

**Assessment of alterations in neurometabolites and brain
microstructure after repetitive mild traumatic brain injury and its
association with the behavioral outcome**

Ph.D THESIS

*Submitted in partial fulfilment of the
requirements for the award of the degree of*

DOCTOR OF PHILOSOPHY

in

BIOTECHNOLOGY

by

MEGHA KUMARI



**DEPARTMENT OF BIOTECHNOLOGY
DELHI TECHNOLOGICAL UNIVERSITY**

DECEMBER 2024

*Dedicated To My Family,
Teachers and Friends*

CERTIFICATE

This is to certify that the thesis entitled, "**Assessment of alterations in neurometabolites and brain microstructure after repetitive mild traumatic brain injury and its association with the behavioral outcome**" being submitted by Ms. Megha Kumari, 2K18/PHDBT/15, Department of Biotechnology, Delhi Technological University, Delhi is a record of bonafide research work carried out by her. Ms. Megha kumari has worked under our guidance and supervision. She has fulfilled the requirements for the submission of this thesis, which to our knowledge has reached the requisite standard.

Co- supervisor

Dr. Richa Trivedi

Scientist 'F'

Institute of Nuclear Medicine and Allied Sciences

DRDO

Delhi - 110054

Supervisor

Prof. Yasha Hasija

Professor

Department of Biotechnology

Delhi Technological University

Delhi-110042

DECLARATION

I, **Megha Kumari**, hereby declare that the thesis, entitled **“Assessment of alterations in neurometabolites and brain microstructure after repetitive mild traumatic brain injury and its association with the behavioral outcome”**, submitted to the Delhi Technological University, in partial fulfilment of the requirements for the award of the Degree of Doctor of Philosophy in **Biotechnology** is a record of original and independent research work done by me during **October 2017 –May 2024** under the Supervision and Guidance of Dr. **Richa Trivedi** Department of **NMR Research Centre, Institute of Nuclear Medicine and Allied Sciences (INMAS), Delhi - 110054** and it has not formed the basis for the award of any Degree / Diploma / Associateship / Fellowship or other similar title to any candidate in any University.

Signature of the Candidate

ACKNOWLEDGEMENT

I would like to express my sincere gratitude to my supervisors **Prof. Yasha Hasija** and my co- supervisor **Dr. Richa Trivedi**. Their unwavering support, insightful suggestions, boundless patience, endless motivation, and expert guidance have been instrumental in my growth as a research scholar. This journey would have been impossible without their contribution. Their support was an unyielding force that carried me through the most challenging times of my life.

My Heartfelt thanks to the **present Director of INMAS**, Dr Sudhir Chandna, for providing me with all the necessary facilities for my research work. I would also like to thank **past Directors of INMAS**, Dr A.K. Singh, Dr Tarun Sekhri, and Dr A.K. Mishra, for their support and encouragement. Bountiful thanks to Dr Vijayakumar Chinnadurai, Dr Poonam Rana, Dr Sonia Gandhi, Dr B.S Hemanth Kumar, Dr Shilpi Modi and Dr Priyanka Sharma for their generous help.

I want to extend my deepest gratitude to all the senior members, Dr. Kavita Singh, Dr Sushant Kumar Mishra, Dr Mukesh Kumar Saini, Dr Ruchi Baghel, and Dr Mamta Gupta and fellow lab mates and friends Dr.Aleem, Dr. Isha Gupta, Deepak Sharma, Abhinit Kumar, Meenakshi Bhardwaj, Dr Kiran Mann, Seema Dhariwal, and Palkin Arora from INMAS, for their kind support and endless scientific interactions in completing my research work.

I am also thankful to the technical staff, Apoorva Sharma and Gautam Sir from INMAS. Mr. Chhail Bihari Singh and Mr. Jitender Singh, from DTU for their assistance and support while conducting my experiments.

I would like to thank the University Grant Commission (UGC) for funding that helped me conduct PhD research.

I owe a debt of gratitude to **my family members** for their love, support and confidence. I am immensely grateful to my parents, Mr. Muktinath Mukhiya and Ms. Prabha Kumari, for being my guiding light and constant support throughout my academic journey. I am fortunate to have wonderful sisters, Shweta and Ruchika, who helped me deal with challenging situations by showering their unconditional love.

Lastly, I would like to thank everyone I may have forgotten to include here for them invaluable contribution. Last but not least, I am sincerely grateful to **God**, the Almighty for instilling in me the idea of exploration as a way of life and bestowing me with ample opportunities, patience, positivity, courage, and wisdom to proceed in this journey

Megha Kumari

ABSTRACT

Traumatic brain injury (TBI) is a leading cause of mortality and morbidity worldwide, affecting both civilian populations and military personnel. Survivors of TBI often experience long-term, disabling changes in cognition, motor function, and personality. The Glasgow Coma Scale (GCS) is a commonly used tool for assessing the level of consciousness and severity of TBI, evaluating three key components: eye-opening response, motor response, and verbal response. Based on the score, TBI is classified into three categories: mild, moderate, or severe. In addition to the GCS, imaging techniques such as CT and MRI scans are often used to assess the extent of injury, particularly in cases of mild TBI. TBI is inherently heterogeneous, meaning it can manifest in various ways depending on the individual. To model this complexity in humans, preclinical studies have used experimental models that replicate the key pathophysiological features of different types of human TBI. Despite the identification of numerous promising neuroprotective agents in experimental studies, none have led to significant improvements in long-term clinical outcomes. Several factors contribute to these translational failures, including differences in the types of data collected (such as histopathological, behavioral, and imaging data) and the timing of data collection (ranging from hours to weeks post-TBI) between clinical and experimental studies. Additionally, preclinical testing should involve multiple experimental TBI models, preferably across different species, to better simulate human conditions.

Repetitive mild traumatic brain injury (rmTBI) refers to a condition where a person sustains multiple mild TBIs over time. Athletes in contact sports like football, hockey, and soccer, military personnel exposed to blasts, and individuals in high-risk activities for falls or head impacts are all vulnerable to rmTBI. The cumulative effects of repeated mild TBIs mean that each successive injury may have a more significant impact on brain function and recovery. Symptoms of mild TBI, such as headaches, dizziness, memory issues, concentration problems, mood changes, or fatigue, can be subtle and temporary, making detection difficult. While each mild TBI may heal relatively quickly, when injuries occur consecutively, recovery can become delayed or incomplete. Some individuals may experience persistent post-concussive symptoms or develop long-term issues. Repeated mild TBIs also increase the likelihood of subsequent injuries and lower the threshold for more severe symptoms. This cumulative effect raises

concerns about long-term consequences, such as chronic traumatic encephalopathy (CTE), a degenerative brain disease linked to repeated brain trauma. Metabolomics, the study of metabolites, offers a way to identify biomarkers associated with TBI. These biomarkers can be used to predict the severity, progression, and recovery of the injury. Nuclear magnetic resonance (NMR) metabolomics plays a vital role in TBI research, helping to understand the metabolic changes that occur following injury. In this study, NMR metabolomics was used to standardize the classification of diffuse TBI in rats and assess its impact on both the brain and serum. Two types of injury models were used: (1) blunt trauma using the modified Marmarou's weight drop model, and (2) blast trauma using a compression-driven shock wave tube.

Chapter 1 provides an overview of TBI, the history of NMR spectroscopy, its basic principles, and its application in metabolomics and TBI research. It also discusses various injury models, including those for repetitive TBI.

Chapter 2 explores graded traumatic brain injury (TBI) in animal models, which simulate varying levels of injury severity, from mild to moderate to severe. This chapter offers valuable insights into the pathophysiological mechanisms of diffuse graded TBI and its effect on the metabolic homogeneity of the hippocampus at acute time points, using ^1H -NMR metabolomics.

Chapter 3 focuses on the differential effects of diffuse brain injury on regions distant from the impact site, such as the hippocampus, thalamus, and striatum, at acute, early sub-acute, and sub-acute time points. Using ^1H -NMR metabolomics, this chapter aims to deepen our understanding of the metabolic changes and pathophysiology of TBI in these brain regions.

Chapter 4 examines the metabolic changes associated with repetitive blast concussions, specifically in female rats. This study investigates how repetitive blast injury reprograms metabolism and explores the interplay between metabolic changes and epigenetic alterations following such injuries.

Chapter 5 investigates TBI-induced metabolic alterations over time, from acute to chronic stages, following both mild and repetitive mild TBI. Using ^1H -NMR metabolomics to analyze serum metabolic

changes, the study also employs fecal 16S rRNA sequencing to explore changes in the gut microbiome following injury.

Chapter 6 summarizes the behavioral changes observed in animals subjected to two types of TBI: diffuse blunt injury and blast-induced injury. This chapter highlights significant behavioral differences between the two injury groups, particularly at chronic time points following the initial injury.

Chapter 7 provides a comprehensive summary of the thesis findings, offering key insights into the pathophysiological and behavioral changes induced by blast and blunt TBI over time.

Table of Contents

• Certificate	i
• Declaration.....	ii
• Acknowledgements.....	iii- iv
• Abstract.....	v-vii
• Table of Contents.....	viii-xi
• List of Figures and table	xii-xiii
• Glossary of Symbols and Abbreviations.....	xiv-xvii
1. Chapter 1	1-23
1.1. Introduction and review of literature.....	1-2
1.2. Primary injury	3
1.3. Secondary injury cascades.....	3-6
1.4. Mechanisms of blast and blunt-induced brain injury.....	7-8
1.4.1. Mechanisms of blast-induced TBI.....	7
1.4.2. Mechanisms of Blunt-induced TBI.....	8
1.5. Animal models of TBI	9
1.5.1. Fluid percussion injury models.....	9
1.5.2. Controlled cortical impact injury model.....	9
1.5.3. Weight drop TBI model.....	10
1.5.4. Marmarou weight drop apparatus.....	10
1.5.5. Blast induced TBI model	11-12
1.6. Metabolomics in TBI	13-14
1.7. Histological and neurobehavioral changes in experimental TBI.....	14-15
1.8. Repeated TBI (Traumatic Brain Injury)	16-18
1.9. Brief history of NMR.....	18-19
1.9.1. Physical basis of NMR	19-22
1.10. NMR metabolomics in TBI	22-23
2. Chapter 2.....	24-35
2.1. Introduction.....	24-26
2.2. Materials and Method.....	26-29
2.2.1. Chemicals	26

2.2.2. Study Model	26
2.2.3. Injury Model	26
2.2.4. Probit Analysis	27
2.2.5. Sample collection and processing	28
2.2.6. NMR data acquisition and processing / Histological staining ...	28-29
2.3. Statistical analysis.....	29
2.4. Result.....	29-
2.4.1. Metabolic changes in hippocampal tissues after graded injury....	30-31
2.5. Discussion.....	32-35
2.6. Conclusion.....	35
3. Chapter 3.....	35 -53
3.1. Introduction.....	35-38
3.2. Materials and Method.....	39
3.2.1. Chemicals.....	39
3.2.2. Study Model.....	39
3.2.3. Injury Model.....	39
3.2.4. Sample collection and processing.....	40
3.2.5. NMR data acquisition and processing / Histological staining.....	40-41
3.3. Statistical analysis.....	41
3.4. Result.....	41-44
3.4.1. Change in metabolic levels among different brain regions in control.....	42
3.4.2. Temporal changes in metabolites within a region after injury.....	42-43
3.4.3. Pathway analysis.....	43-45
3.5. Discussion.....	46-52
3.6. Conclusion	53
4. Chapter 4	55-66
4.1. Introduction.....	55- 56
4.2. Materials and	
Method.....	56-57
4.2.1. Chemicals.....	56-57
4.2.2. Study design.....	57

4.3. Blast exposure.....	57-58
4.3.1. Immunohistochemistry analyses.....	58
4.4. Sample preparation for NMR and spectra acquisition.....	58-59
4.5. Concentration of metabolites	59-60
4.6. Statistical analysis	60
4.7. Results.....	60-63
4.7.1. Neurometabolic alteration among the groups	61
4.7.2. Correlation among the neurometabolites	62
4.7.3. Histological changes among the groups	62-63
4.8. Discussion.....	64-65
4.9. Conclusion.....	66
5. Chapter 5.....	67-78
5.1. Introduction	67-69
5.2. Material and method	69-70
5.2.1. Chemicals	69
5.2.2. Animal model	69
5.2.3. Sample collection and preparation	69-70
5.3. NMR spectra acquisition	70
5.4. Fecal 16S rRNA sequencing.....	70-71
5.5. Statistical analysis.....	72
5.6. Results	72-74
5.7. Discussion	75-
5.7.1. Systemic change in serum metabolome following diffused closed head injury	75-77
5.7.2. Gut microbiome changes after mild and rmTBI	77-78
5.8. Conclusion	78
6. Chapter – 6.....	79-89
6.1. Introduction	79-80
6.2. Materials.....	80
6.3. Method	80-81
6.3.1. Blunt trauma.....	81
6.3.2. Blast exposure.....	81
6.4. Behavioral paradigm	82-83
6.4.1. The Neurobehavioral Severity Scale (NSS)	82
6.4.2. Rotating pole	82
6.4.3. Open field Test	82-83
6.4.4. Novel object Recognition Test (NORT)	83
6.5. Statistical analysis	83
6.6. Results	83
6.6.1. Behavioral changes after repetitive TBI	83-87

6.7. Discussion.....	87
6.7.1. Alteration in NSS after Blast and blunt TBI	87-88
6.7.2. Alteration in rotating pole after blast and blunt TBI	88
6.7.3. Alteration in cognitive function after blast and blunt TBI	88-89
6.8. Conclusion	89
7. Chapter 7	90-95
7.1. Summary and conclusion	90-95
8. Reference	96-116.
9. Publication	117-118

List of figures and table

Figure1.1: Marmarou Weight Device	6
Figure1.2: compression driven shock wave tube.	7
Figure 2.1: Survival curve.	27
Figure 2.2: Representative NMR spectra with selected metabolites	30
Figure 2.3 Showing metabolic change in hippocampal tissue after 24hrs of injury.	31
Figure 2.4: Showing the presence of neurite plaques in hippocampus after 24hrs	
of injury.	32
Figure 3.1: Line graph of all selected metabolites in hippocampus, thalamus	
and striatum at all three time-points. P- Value < 0.05, considered significant.	23
Figure 4.1: Graph showing the significantly changed hippocampal tissue metabolites after	
brain injury.	32
Figure 4.2: Heat diagram of correlation analysis between metabolites.	33
Figure 4.3: GFAP positive cells in the dentate gyrus region of hippocampus of different	
injury groups and bar graph of GFAP positive cells at 24hrs PI.	33
Figure 4.4: AH3 positive cells in the dentate gyrus region of hippocampus	
of different injury groups and bar graph of AH3 positive cells at 24hrs PI.	33
Figure 5.1a, 5.1b, and 5.1c: showing the change in gut microbiota after 30days	

of injury.	38
Figure 5.2: Graph showing the significantly changed serum metabolites after blunt TBI.	39
Figure 6.1: Showing graph of NSSR experiment following blunt TBI.	45
Figure 6.2: Showing graph of NSSR experiment following blast TBI.	45
Figure 6.3: Line graph of rotating pole experiment following blunt TBI.	45
Figure 6.4: Line graph of rotating pole experiment following blast TBI.	46
Figure 6.5: Showing the NORT and OFT results following blunt TBI.	46
Figure 6.6: Showing the NORT and OFT results following blast TBI.	47
Table1: List of significantly altered pathway following moderate injury.	24
Supplementary Table1: List of selected metabolites and their peak details.	54
Supplementary Table2: Concentration of metabolites.	68

GLOSSARY OF SYMBOLS AND ABBREVIATIONS

°C	degree centigrade
1H NMR	Proton Nuclear magnetic resonance
1H-MRS	proton magnetic resonance spectroscopy
ACHI	awake-closed head injury
AD	axial diffusivity
ADP	Adenine diphosphate
AH3	acetyl histone 3
alpha-KG	alpha-ketoglutarate
AMPARs	α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors
ANOVA	Analysis of variance
APP	amyloid precursor protein
ATP	Adenosine triphosphate
BBB	blood-brain barrier
BCA	branched-chain amino acids
blTBI	Blast induced traumatic brain injury
CCI	Controlled cortical impact
CN	Composite Neuroscore
CNS	central nervous system
COSY	Correlation Spectroscopy
CPMG	Carr-Purcell-Meiboom-Gill
CTE	chronic traumatic encephalopathy
DAI	Diffuse axonal injury
DNA	Deoxyribonucleic acid
DTI	Diffusion tensor imaging
EPM	Elevated Plus Maze

FA	fractional anisotropy
FAD	flavin adenine dinucleotide
FID	Free induction decay
FPI	Fluid Percussion Injury
FT	Fourier transform
g	gram
GABA	gamma-aminobutyric acid
GFAP	glial fibrillary acidic protein
h	hour
h	hour
HMDB	Human metabolome database
HPLC	High-Performance Liquid Chromatography
Iba1	ionized calcium-binding adapter molecule 1
IHC	Immunohistochemical
IL	interleukin
LDL	low-density lipoprotein
MBP	Myelin binding protein
MI	Myo inositol
min	minute
mL	milliliter
moTBI	moderate traumatic brain injury
mTBI	mild traumatic brain injury
MWM	Morris Water Maze
NAA	N-acetyl aspartic acid
NAD	Nicotinamide adenine dinucleotide
NADP+	Nicotinamide adenine dinucleotide phosphate
NHE 1	Na ⁺ /H ⁺ exchanger protein

NMDAs	N-methyl-D-aspartate receptors
NOESYPR	Nuclear overhauser effect spectroscopy presaturation
NORT	Novel Object Recognition Test
NSS	Neurobehavioral Severity Scale
OFT	Open Field Test
PBS	Phosphate buffered saline
PC	phosphatidylcholine
PCA	Principle component analysis
PCs	Principle components
PE	phosphoethanolamine,
PFA	para formaldehyde
PI	post injury
PLS-DA	Partial least squares discriminant analysis
PPM	parts per million
Q2	predictive ability
R2	goodness of fit
RF	radiofrequency
rmTBI	repetitive traumatic brain injury
ROS	Reactive oxygen species
rpm	revolutions per minute
s	second
SD	Standard deviation
T1	longitudinal relaxation
T2	transverse relaxation
TBI	traumatic brain injury
TCA	Tricarboxylic acid
TLR	Toll-like receptor

TSP	Trimethylsilyl-2,2,3,3-tetra-deuteropropionic acid
v/v	volume/volume percentage
VIP	Variable importance in projection
VLDL	Very Low-Density Lipoprotein
ZGPR	Zero-Go-Pulse-Relaxation
Δ	Chemical Shift
μL	microliter
μm	micrometer
kPa	kilopascal
GCS	Glasgow Coma Scale
CT	Computed Tomography

Chapter 1

Introduction and Review of literature

1.1 Introduction

Traumatic brain injury (TBI) is a major public health concern, responsible for one-third of all injury-related fatalities (Roozenbeek et al., 2013). Worldwide, TBI affects over 27 million individuals, with motor vehicle accidents, falls, and assaults being the leading causes of hospitalization (James et al., 2019). It is a primary contributor to both mortality and long-term disability, affecting civilian and military populations alike. Those who survive often face persistent cognitive, motor, and personality changes, highlighting the urgent need for further research in this area. TBI is defined as a disruption in brain function or pathology caused by an external force (Menon et al., 2015). This broad definition encompasses a complex range of injury types, which often overlap, and recent efforts have focused on distinguishing these subtypes to develop more precise treatments based on specific pathobiological processes (Strianese et al., 2020). These approaches must also account for variations in host responses, which may be influenced by concurrent trauma or pre-existing conditions. TBI involves both primary and secondary injury mechanisms. Primary injury occurs immediately upon impact, resulting in mechanical damage to brain tissue such as contusions, hemorrhage, and axonal shearing. Secondary injury, which develops in the minutes to months following the initial trauma, involves a cascade of cellular and molecular events that cause further brain cell death, tissue damage, and atrophy. Unlike primary injury, secondary injury provides a window for therapeutic intervention, offering opportunities to prevent additional damage and improve long-term recovery. TBI severity is typically classified as mild, moderate, or severe. The Glasgow Coma Scale (GCS) is commonly used to assess TBI severity. However, the evolving nature of secondary injury and its associated pathophysiological responses highlight the complexity of TBI as a disease, which complicates treatment and patient care (Masel & DeWitt, 2010). Mild, moderate, and severe TBI are classified according to GCS scores, with mild TBI

ranging from 13 to 15, moderate TBI from 9 to 13, and severe TBI from 3 to 8. Even mild to moderate injuries can result in cognitive and behavioral impairments. The rising incidence of Alzheimer's disease, chronic traumatic encephalopathy (CTE), and Parkinson's disease in individuals with severe or repeated mild TBI has drawn increasing concern. Mild TBI, the most common form, can cause neurological, cognitive, and behavioral changes. Neurological symptoms may include headaches, nausea, dizziness, sleep disturbances, and vision issues, while cognitive symptoms can involve attention, memory, language, communication, and executive function impairments. Behavioral changes often include irritability, aggression, impulsiveness, and poor social judgment.

Animal models are essential for understanding the mechanistic changes associated with TBI, as they help simulate clinical scenarios and produce reproducible results. Clinical TBI cases exhibit varying pathophysiological conditions due to factors such as injury location, severity, and pre-existing conditions, including genetics, age, health, gender, and substance abuse. In India, limited data indicates that TBI occurs frequently, with a notable percentage of cases (69%) occurring in individuals aged 15-35 years (Verma & Tewari, 2004). Road traffic accidents (60%), falls (20-25%), and violence (10%) are the leading causes of TBI (Gururaj, 2002). Developing countries like India suffer significant socioeconomic losses due to TBI, and the country has the highest global fatality rate from road traffic accidents (Samabasivan, 1991). Approximately 70% of TBI cases are classified as mild (mTBI) (Belanger, 1996), with 20-30% falling into the moderate or severe categories (moTBI). Alarmingly, about 40% of individuals with mTBI do not seek medical care (Sosin, 1991). Mild TBI can lead to postconcussion symptoms, which include neurological, cognitive, and behavioral manifestations.

Although most of these symptoms resolve within 10 days to two weeks post-injury (Meehan et al, 2011), more than 25% of individuals continue to experience symptoms long after the injury (Dikmen, 2010; Lannsjo, 2009; Sigurdardottir, 2009).

1.2 Primary Injury

Primary injury in traumatic brain injury (TBI) refers to the initial damage due to direct impact to the head, rapid acceleration or deceleration, penetrating injuries, or blast events. This damage often results in membrane disruption, axonal injury, and blood-brain barrier (BBB) breakdown. Diffuse axonal injury (DAI) is one of the most prevalent pathologies in closed head injuries, often leading to coma (Smith et al., 2000). Primary axotomy occurs when axons are directly severed during TBI, leading to cytoskeletal distortion (Povlishock & Christman, 1995). Additionally, micropores may form in cell membranes, allowing sodium and calcium influx, which causes depolarization of the membrane potential (Pettus et al., 1994; Farkas & Povlishock, 2007; Krishnamurthy & Laskowitz, 2016). The blood-brain barrier (BBB), is a structure which separates brain parenchyma from the blood, consists of endothelial cells connected by tight junctions. This barrier regulates the influx of nutrients and ions while restricting harmful molecules, ensuring homeostasis and maintaining an immunologically privileged environment. TBI disrupts the BBB by damaging endothelial cells, disrupting tight junctions, and permitting atypical molecular transport (Campbell et al., 2012). This disruption contributes to neuronal dysfunction and initiates secondary injury cascades, compounding brain damage.

1.3 Secondary Injury Cascades

Secondary injury encompasses the pathological processes triggered by the primary injury. The central mechanism of secondary injury is excitotoxicity (Krishnamurthy & Laskowitz, 2016), characterized by an excessive release of excitatory neurotransmitters, particularly glutamate, leading to calcium overload in post-synaptic neurons. Under normal conditions, glutamate released from pre-synaptic neurons into the synaptic cleft binds to ligand-gated ion receptors such as N-methyl-D-aspartate receptors (NMDARs) and α -amino-3-hydroxy-5-methyl-4isoxazolepropionic acid receptors

(AMPARs) on post-synaptic membranes, facilitating sodium and calcium influx. However, after TBI, excessive glutamate release occurs in proportion to injury severity, with extracellular glutamate levels rising up to 50-fold compared to normal levels (Faden et al., 1989; Palmer et al., 1994). This abnormal glutamate surge overwhelms the system, impairing the function of astrocytic glutamate transporters and exacerbating calcium influx into neurons. This cascade disrupts normal neuronal signaling, contributes to cellular damage, and perpetuates the cycle of neurotoxicity, ultimately leading to widespread neuronal death and further brain dysfunction. Increased intracellular calcium can lead to further neuronal damage and even death by several means, one of which is mitochondrial dysfunction. Mitochondria have been described as “calcium sinks” as they will sequester large amounts of calcium from the cytosol in an effort to maintain calcium homeostasis (Ichas and Mazat 1998). Under injury conditions where calcium is overloaded in the cell, the large quantities of calcium taken up by the mitochondria lead to increases in mitochondrial production of reactive oxygen species (ROS) (Sullivan, Thompson et al. 1999; Chen, Yang et al. 2007), loss of ATP production (Jacobus, Tiozzo et al. 1975), and opening of the mitochondrial permeability transition pore (Bernardi 1996). The consequence of these events is the release of cytochrome c and other pro-apoptotic factors which trigger the formation of the caspase-3 cleaving apoptosome. Cleavage of caspase-3 induces apoptosis of the cell, resulting in cell death.

Another feature of calcium overload after TBI is activation of cytoskeletal-cleaving calpains (Araujo, Carreira et al. 2010). Breakdown of filaments and microtubules can cause damage to the structural integrity of the neuron (Liu, Liu et al. 2008) and dysfunction of the transported networks within the neuron (Saatman, Abai et al. 2003). Immunohistochemical labeling of transport proteins after TBI, such as amyloid precursor protein (APP), have been used to identify damaged axons within hours (Roberts, Gentleman et al. 1994). However, not all axons within a region appear damaged by the primary injury, likely due to the heterogeneity among axons in their size and myelination. In contrast, some dysfunctional axons can also appear normal depending on the marker used to detect injury (Tomei,

Spagnoli et al. 1990). Calpains are also responsible for cleaving the mGluR1 receptor blocking the neuroprotective PI3K-AKT pathway (Liu, Liu et al. 2008). These events are spread via positive feedback, as the initial glutamate release in injured neurons induces further neurotransmitter release from the postsynaptic cell (Werner and Engelhard 2007). Disruption of the BBB initiates additional secondary injury cascades by allowing an influx of thrombin, fibrinogen and albumin into the parenchyma initiating a coagulation cascade (Price, Wilson et al. 2016). Decreased cerebral blood flow causes ischemia at the site of impact. A lack of oxygen and nutrients in the impact area limits the production of ATP needed for cell function (del Zoppo and Mabuchi 2003; Schwarzmaier, Kim et al. 2010). Alongside coagulation factors, neutrophils and peripheral immune cells, such as macrophages, also infiltrate brain tissue (Chodobska, Zink et al. 2011). Because the brain is considered immunologically privileged, the infiltrating immune cells are believed to become active, secreting cytokines (Kumar and Loane 2012). Secretion of these cytokines then initiates the activation of microglia and astrocytes. Microglia and astrocytes express pattern recognition receptors on their cell surface called Toll-like receptors (TLR) that allow them to recognize foreign or damaged cells (Gorina, Font-Nieves et al. 2011). Recognition by TLRs of trigger molecules induces the production and release of inflammatory chemokine and cytokines initiating the inflammatory cascade. Under normal conditions, astrocytes play an important role in maintaining homeostasis of neurons by the uptake of glutamate from synapses (Chen and Swanson 2003). Under injury conditions in which excessive amounts of neurotransmitters are released into the synaptic cleft, astrocytes downregulate production of glutamate transporters (Beschorner, Simon et al. 2007). This down-regulation is believed to perpetuate the excessive amounts of glutamate in the synaptic cleft resulting in cell death. Astrocytes are also responsible for maintaining the integrity of the BBB (Risau and Wolburg 1990). Following TBI, astrocytes secrete inflammatory molecules that modulate reactivity and suppression of astrocyte function (Gorina, Font-Nieves et al. 2011). Secretion of the cytokine IL-6 by astrocytes has been reported to increase the permeability of the BBB (Schwaninger, Sallmann et al. 1999), further damaging

the already delicate BBB allowing for greater infiltration of peripheral cells (Jin, Liu et al. 2012). In addition, astrocytes increase expression of glial fibrillary acidic protein (GFAP) and vimentin, two intermediate filament proteins (Pekny, Eliasson et al. 1999 Intermediate Filament Protein Partnership in Astrocytes and increase self-proliferation in areas of injury (Bardehle, Kruger et al. 2013). Within a week following injury in rodent models, reactive astrocytes have been shown to form a glial scar which effectively surrounds the injured tissue separating it from nearby healthier tissue (Villapol, Byrnes et al. 2014). The glial scar is beneficial in stopping the influx of peripheral immune cells that contribute to a continued inflammatory response (Dardiotis, Hadjigeorgiou et al. 2008), but creates a barrier for the regeneration of damaged axons. Microglia are the brain's resident immune cells. In their resting state, they constantly monitor the brain's environment, looking for signs of damage or infection. When injury occurs, microglia become activated, releasing pro-inflammatory cytokines to enhance their reactivity and assist in clearing debris through phagocytosis. While the removal of debris is beneficial there is growing literature that microglia can remain activated long after injury and may contribute to continued neurodegeneration (Block and Hong 2005; Chen, Johnson et al. 2009).

Similar to astrocytes, microglia exhibit heterogeneity. Activated microglia are thought to exist in three distinct states: classically activated, alternatively activated, and acquired deactivation. Classically activated microglia (also referred to as M1 stage) are believed to release proinflammatory cytokines and reactive oxygen species, which can lead to white matter damage and cell death (Karve, Taylor et al., 2016). In contrast, alternatively activated microglia (M2b stage) are thought to play a role in tissue repair and healing by producing scavenger receptors. Microglia in acquired deactivation states (also known as stage M2c) promote inflammation resolution (Cao, Thomas et al. 2012; Karve, Taylor et al. 2016). Several studies have reported a transient increase in M2 microglial responses following injury that is followed by a delayed increase in M1 microglia (Hu, Li et al. 2012; Wang, Zhang et al. 2013) suggesting the activated microglia observed chronically after injury are deleterious.

1.4 Mechanisms of Blast and Blunt-Induced Traumatic Brain Injury

The mechanisms of blast-induced and blunt-induced traumatic brain injuries involve complex and distinct processes. Blast-induced TBI is primarily caused by rapid pressure changes from blast waves, leading to diffuse axonal injury, neuroinflammation, and oxidative stress. In contrast, blunt-induced TBI results from direct mechanical forces, causing coup-contrecoup injuries, shear forces, and skull fractures. Research into these mechanisms continues to provide valuable insights into the pathophysiology of TBIs and informs the development of targeted treatments to improve outcomes for affected individuals.

Blast-Induced Traumatic Brain Injury (bTBI)

Blast-induced TBI primarily affects military personnel and individuals exposed to explosive devices. The unique nature of blast waves results in complex injury mechanisms that differ from other forms of TBI.

1.4.1 Mechanisms of Blast-Induced TBI:

- **Primary Blast Injury:** This is caused by the direct impact of the blast wave on the brain. The rapid overpressure followed by underpressure can cause significant brain injury without physical penetration. The mechanisms include:
- **Rapid Pressure Changes:** The blast wave induces rapid compression and decompression of brain tissues, leading to shear forces and diffuse axonal injury (DAI) (Bauman, et al., 2009).
- **Cavitation:** The negative phase of the blast wave can create cavities within brain tissues, leading to microvascular damage and hemorrhage (Garman, et al., 2011).
- **Secondary Blast Injury:** This results from shrapnel and debris propelled by the blast. The impact can cause penetrating brain injuries and additional blunt trauma.

- **Tertiary Blast Injury:** The blast wave can displace the body, causing it to collide with objects. This can lead to blunt trauma, similar to other forms of TBI.
- **Quaternary Blast Injury:** This includes all other injuries from the blast, such as burns and inhalation of toxic gases, which can exacerbate brain injury.

Blunt-Induced Traumatic Brain Injury

Blunt-induced TBI typically results from impacts to the head, such as falls, motor vehicle accidents, or sports injuries. The mechanisms of injury involve direct mechanical forces applied to the skull and brain.

1.4.2 Mechanisms of Blunt-Induced TBI:

- **Coup-Contrecoup Injury:** When the head strikes an object, the brain impacts the inside of the skull at the site of impact (coup) and the opposite side (contrecoup). This can cause bruising, bleeding, and axonal damage.
- **Coup Injury:** The brain is damaged directly under the site of impact.
- **Contrecoup Injury:** The brain is damaged on the opposite side due to the rebound effect (Gennarelli, et al., 1982).
- **Shear Forces:** Rotational forces during the impact cause shearing of brain tissues, leading to diffuse axonal injury (DAI) (Adams, et al., 1982).
- **Skull Fractures:** Blunt force can cause fractures of the skull, leading to additional brain damage from bone fragments and increased intracranial pressure.

1.5 Animal models of TBI

Animal models of traumatic brain injury (TBI) play a crucial role in advancing our understanding of TBI pathophysiology, evaluating potential treatments, and exploring therapeutic mechanisms. Animal models allow us to recreate TBI conditions in a controlled and reproducible manner. By inducing controlled injuries that mimic the mechanical forces and impact experienced in human TBI, these models help in understanding pathophysiology, assessing neurobehavioral outcomes, exploring therapeutic interventions, investigating long-term effects and translational research.

1.5.1 Fluid percussion injury models

Fluid Percussion Injury (FPI) is a widely utilized and well-established model for traumatic brain injury (TBI). It has been effectively applied in several animal species, including rabbits, cats, rats, mice, and pigs. The injury can be delivered either centrally (midline FPI), along the sagittal suture between bregma and lambda, or laterally (lateral FPI), over the parietal cortex.

This model produces graded neurological, histological, and cognitive outcomes that closely resemble the effects of human TBI (Dixon, 1987). The injury is induced by delivering a brief fluid pulse (~20 msec) to the dura mater through a craniotomy, causing a temporary deformation of the brain tissue. By varying the intensity of the fluid pulse and the location of the impact, the FPI model can mimic both diffuse and focal injury types, resulting in corresponding neurological impairments and neuropathological alterations. In the FPI model, injury is generated by a pendulum striking a piston connected to a fluid reservoir, which creates a pressure pulse transmitted to the dura mater through the craniotomy. This can be applied centrally along the midline or laterally over the parietal bone, between bregma and lambda. The impact causes a brief displacement and deformation of the brain tissue. The severity of the injury depends on the intensity of the pressure pulse (McIntosh et al., 1989)

1.5.2 Controlled cortical impact injury model

The CCI model uses a pneumatic or electromagnetic impact device to drive a rigid impactor onto the exposed intact dura, and mimics cortical tissue loss, acute subdural hematoma, axonal injury, concussion, blood-brain barrier (BBB) dysfunction and even coma (Morales DM, et al, 2005). The advantage of this injury model over other TBI models is the ease at which mechanical factors, such as time, velocity and depth of impact, can be controlled; thus, it may be more useful than the FPI model for biomechanical studies of TBI (Wang HC, Ma YB., 2010).

1.5.3 Weight drop TBI model

In weight drop models, the skull is exposed (with or without a craniotomy) to a free falling, guided weight (Morales DM, et al, 2005). Injury severity in these models can be altered by adjusting the mass of the weight and the height from which it falls. One such example is

Marmarou weight drop apparatus.

1.5.4 Marmarou Weight Drop Apparatus

One widely used model is Marmarou's weight drop model, which simulates diffuse brain injury in rodents. The Marmarou's weight drop model was developed in the 1990s by Anthony Marmarou to create a reproducible and clinically relevant method for studying diffuse brain injury in animals. This model involves dropping a weight from a predetermined height onto a rodent's head, causing rapid acceleration and deceleration forces similar to those experienced in human TBIs. Marmarou's weight drop model of traumatic brain injury has been a cornerstone in TBI research, providing valuable insights into the pathophysiology and potential treatments for diffuse brain injury. Its ability to replicate key features of human TBIs, such as diffuse axonal injury and neuroinflammation, makes it an indispensable tool for researchers. The animal, typically a rat or mouse, is anesthetized and placed on

a foam bed to absorb the impact. A weight, of 450 grams was dropped from a specified height onto a small metal disc placed on the skull. The impact results in a diffuse brain injury, characterized by widespread axonal damage, edema, and hemorrhage. This model effectively mimics the biomechanical forces of human TBIs, making it a valuable tool for studying injury mechanisms and potential treatments.

Marmarou's weight drop model has significantly contributed to our understanding of TBI's pathophysiology. Its ability to induce diffuse axonal injury (DAI), a common feature in human TBIs, has made it particularly valuable for exploring the cellular and molecular responses to brain injury. Studies have shown that the model induces axonal swelling and disconnection, mirroring DAI observed in humans, neuroinflammation, oxidative stress, blood brain barrier disruption and cognitive and motor deficits (Smith, et al., 1997; Kumar, et al., 2015; Hamm, et al., 1994) .




Figure1.1: Marmarou Weight Device

1.5.5 Blast induced TBI model

Shock wave-generated traumatic brain injury (TBI) models have become a cornerstone in the study of blast-related brain injuries, particularly those experienced by military personnel and victims of explosions. These models aim to replicate the primary blast wave's effects on the brain, providing valuable insights into injury mechanisms and potential therapeutic interventions. Shock wave-

generated TBI models are designed to simulate the primary blast wave, characterized by a rapid rise in pressure followed by a negative phase. This sudden pressure change can cause significant brain injury without physical penetration, making it a crucial area of study for understanding blast-induced TBI. The primary blast wave in shock wave-generated TBI models is typically produced using a shock tube, a device that generates controlled shock waves. The basic setup involves a high-pressure chamber separated from a low-pressure chamber by a diaphragm. When the diaphragm is ruptured, it creates a shock wave that travels through the low-pressure chamber, exposing animal to the blast.

Shock wave-generated TBI models have been instrumental in elucidating the pathophysiological mechanisms underlying blast-induced brain injuries. Previous studies showed these models replicate key features of human bTBI, such as diffuse axonal injury, neuroinflammation, and blood-brain barrier disruption, oxidative stress and cognitive decline (Garman, et al., 2011; Goldstein, et al., 2012; Kuriakose & Younger, 2013; Hoffer, et al., 2013). Another study reported that the intensity and duration of the shock wave significantly influence the severity of brain injury (Taylor & Ford, 2009).




Figure1.2: Compression driven shock wave tube.

1.6 Metabolomics in TBI

TBI pathology and progression appears to have a metabolic basis. Studies have tried to establish evidence of a significant metabolic abnormalities such as oxidative stress (Bayir 2006, Viant 2005), mitochondrial dysfunction and excitotoxicity (Prasad and Bondy 2015, Viant 2005) in TBI. These changes are resultant of alterations in the cellular levels of different metabolites viz glucose, glutamate, lactate, pyruvate, NAA, NAAG, creatines etc (Timofeev 2011). Inflammation has shown to play a key role in TBI pathology (Yatsiv 2002), where metabolites like ATP and lactate have been reported as sensitive markers of inflammation.

These findings suggest that metabolomics could play a pivotal role in investigating TBI (Fitzpatrick and Young 2013). However, there are few studies on TBI metabolomics reported in literature. Earlier study by Viant et al. (2005) employed ^1H NMR to study the brain and plasma metabolome from rats following a TBI model of lateral fluid percussion. They report that the metabolic fingerprint following TBI comprises information on oxidative stress, excitotoxic injury, membrane disruption and neuronal injury (Viant 2005). Pascual (2007) report the use of ^1H MRS to study Sprague–Dawley rats that were subjected to a closed head impact and examined over a period of 72 h. They found that using a stepwise multivariate model they were capable of discriminating amongst the various time groups (1, 9, 24, 48 and 72 h) based on a combination of taurine and myoinositol (Pascual 2007). Casey et al. (2008) reported alterations in cerebral metabolism over a 7 days period of immature rats following TBI. Brain tissue samples were analyzed using ^1H NMR, revealing changes in metabolic processes between 4 and 24 hours after traumatic brain injury (TBI). Specifically, there was an increase in glycolysis and/or oxidative metabolism, as indicated by the elevated ratio of lactate to creatine. Additionally, at 24 hours and 7 days post-TBI, signs of neuronal damage were detected, as reflected by the reduced ratio of N-acetyl aspartate to creatine. These findings suggest that metabolic disturbances begin as early as 24 hours after TBI and persist for up to one week (Casey, 2008). Similarly, Lemaire et al; 2011 used ^1H NMR to study TBI in rats at 3, 24, and 48 hours after injury, noting that lactate, valine, and ascorbate

were the first metabolites significantly altered following TBI. Robertson (2013), building on Casey's study, utilized both ¹H and ¹³C NMR to track the utilization of labeled glucose in the brains of immature rats 24 hours post-TBI. They found that oxidative glucose metabolism in the immature rat brain begins to recover within 24 hours after injury, though they suggested further research to explore the pathways leading to decreased metabolism after TBI. The search for reliable serum biomarkers for TBI has been ongoing, with great potential for enhancing diagnosis, understanding the mechanisms of injury, monitoring disease progression, identifying complications, predicting outcomes, and discovering new treatments (Mondello and Hayes, 2015). The use of NMR spectroscopy to identify brain biomarkers in TBI models, such as in a closed head injury mouse model (Bahado-Singh, 2016), has also been explored. Given that TBI damage progresses over time, serial measurements of metabolite levels are essential for tracking the extent of injury, monitoring therapeutic responses, and making accurate prognoses. Blood-based biomarkers offer a distinct advantage, as they can be repeatedly measured at relatively low cost and effort, which is why they are the focus of significant research interest (Neher, 2014). In our study, we employed NMR-based metabolomics to identify potential serum and urine biomarkers for various severities of TBI.

1.7 Histological and Neurobehavioral changes in experimental TBI

Experimental models of traumatic brain injury (TBI) are crucial for understanding the neurological, cognitive, and pathological changes that occur in human patients. Komoltsev et al. (2013) utilized a lateral fluid percussion injury (FPI) model in rodents to investigate the effects of TBI on behavior and neuronal health. Their findings revealed early memory deficits, observed through new object recognition tests (NORT), linked to elevated corticosterone levels on day 3 post-injury. Rats also displayed mild behavioral impairments, including delayed spatial memory deficits in the Barnes maze, which persisted for up to three months. Changes in stress reactivity were noted, including a reduced

corticosterone response to restraint stress and altered reactions to forced swim stress. Additionally, TBI resulted in astrogliosis, including increased GFAP staining, which suggested activated astrocytes and potential morphological alterations observed three months after injury (Komoltsev et al., 2013). In a study by Hylin et al. (2013), mild TBI (mTBI) was induced using 1.0 and 1.5 atm overpressure FPI. Both injury intensities led to acute, transient suppression of neurological functions without visible brain contusion. However, the 1.5 atm injury caused temporary motor disturbances and significant impairments in spatial learning and short-term memory. This injury also resulted in a marked reduction in cerebral perfusion at the injury site, accompanied by a robust inflammatory response, evidenced by increased GFAP and Iba1 immunoreactivity in the corpus callosum and thalamus. Further, a significant reduction in fractional anisotropy (FA) values in the cingulum, along with increased silver impregnation, indicated axonal injury in this region as well as in the corpus callosum and internal and external capsules.

In a study using a controlled cortical impact (CCI) model, rats showed higher mean kurtosis and mean diffusivity values in the ipsilateral perilesional cortex and hippocampus, with decreased FA values in the corpus callosum. This was accompanied by increased GFAP and Iba1 staining, and reduced NeuN and MBP staining in all ipsilateral regions, suggesting considerable neural damage. No significant differences were observed in the contralateral regions for diffusional kurtosis imaging parameters or immunohistochemical staining. Behavioral tests, such as the Morris water maze, revealed longer platform crossing times in the probe test, indicating deficits in cognitive function (Wang et al., 2018). Johnstone (2015) induced moderate lateral FPI in adult male Sprague-Dawley rats and conducted behavioral assessments, MRI, and electrophysiological recordings from the barrel cortex 12 weeks postTBI. The results showed sensorimotor deficits, cognitive impairments, and anxiety-like behaviors, alongside significant atrophy in the barrel cortex and other brain regions. Diffusion tensor imaging (DTI) revealed increased FA, axial diffusivity (AD), and tract density, suggesting long-term recovery of neuronal responsiveness through structural reorganization. Stemper (2015) used a rodent model of

rotational acceleration-induced mTBI to examine the effects of injury magnitude and duration on behavioral and neuroimaging outcomes. While no significant locomotor or cognitive deficits were observed in the Composite Neuroscore (CN) and Morris Water Maze (MWM), the Elevated Plus Maze (EPM) test showed that increased injury duration led to more significant activity and exploratory behaviors. Ex-vivo DTI analysis indicated that changes in FA in the amygdala were associated with both the magnitude and duration of the injury, with extended injury duration resulting in FA changes at the interface between gray and white matter

1.8 Repeated Traumatic Brain Injury

A history of traumatic brain injury (TBI) significantly increases the likelihood of sustaining subsequent head injuries, with individuals being 2 to 5 times more likely to experience another TBI (Guskiewicz et al., 2007; Emery et al., 2011). This heightened risk is especially concerning for those involved in activities such as military training and sports, where repeated head impacts are common (Guskiewicz et al., 2000). However, accurately estimating the incidence of TBIs in these activities is challenging due to two main factors. First, many head injuries go unreported (McCrea et al., 2004), and second, current data reporting systems often categorize TBIs from sports or recreational activities under broader categories like "falls" or "struck by/against" (CDC, 2015). It is widely believed that failing to allow sufficient recovery time between head injuries may result in more severe and lasting pathological and behavioral outcomes. Individuals who have experienced multiple mild TBIs tend to perform worse on cognitive tasks compared to those who have only had a single mild TBI (Gronwall & Wrightson, 1975). A particularly severe consequence of multiple head injuries is second impact syndrome (SIS), where a second head injury occurs before the symptoms from the first injury have fully resolved, leading to dangerously high intracranial pressure and, often, significant morbidity or mortality (McCrory & Berkovic, 1998). While most individuals with mild TBI recover within a week (McCrory et al., 2005; Marar et al., 2012), repeated injuries can lead to more complicated outcomes.

A study by Allen et al. (2023) investigated mild and repetitive mild TBI (rmTBI) in rats and found persistent behavioral deficits in a beam task 1 and 4 hours post-injury. The researchers observed changes in hippocampal metabolism, including increased glutamine and decreased glucose, in rats with repetitive mild TBI, while no significant effects were seen in the single mTBI group. Although there were no significant changes in lipid levels or mitochondrial function in the hippocampus, some alterations in lipids were detected in the cortex following repeated injuries. The study used an awake-closed head injury (ACHI) paradigm to model mild TBI and assessed metabolic changes in the hippocampus 24 hours after the final injury using in vivo proton magnetic resonance spectroscopy (¹H-MRS). Mitochondrial bioenergetics were also measured 30 hours post-injury. Lipidomic evaluations of the hippocampus and cortex were conducted, revealing potential metabolic shifts following repetitive injuries (Allen et al., 2023).

In another study on mild and repetitive mild TBI using the fluid percussion injury (FPI) model, significant changes in righting times were observed in all injury groups compared to sham rats 1 and 2 days after injury. Behavioral testing in the beam walking task revealed notable differences in latencies between groups over a 5-day period. Spatial learning and memory were assessed using the Morris water maze (MWM) task from days 9 to 13 post-injury, with significant differences in spatial learning latencies between the groups and the sham group. Histopathological examination, using Cresyl violet staining, showed cortical thinning, hemorrhaging, and tissue damage in several brain regions, with the most severe damage observed in the qFPI group. Additionally, microglial activity in the cortex and thalamus increased, and changes in the subcortical white matter were detected. GFAP levels were elevated in the subcortical white matter and thalamus, but remained unchanged in the hippocampus and cortex following injury (Fronczak et al., 2022).

Campos-Pires et al. (2023) examined blast-induced unconsciousness in rats and found significant changes in the righting reflex and latency to fall in the accelerating RotaRod test on days 1, 15, and 22 following the blast. Weight loss was observed up to day 22 post-injury. Neuronal loss was found in

specific brain regions, such as the motor cortex, somatosensory cortex, auditory cortex, and amygdala, following repeated blast exposure. However, no significant changes in astrocyte activation were detected in these regions.

Finally, Bielanin et al. (2024) reported on a study of repetitive mild TBI (rmTBI) in a mice model using the controlled cortical impact (CCI) method. They found that rmTBI mice exhibited poorer motor function in the rotarod test at days 10 and 13 post-injury compared to sham mice. Behavioral testing at day 40 post-injury revealed worsened performance in the Ymaze novel spatial recognition test, with significant reductions in differentiation and recognition indices, indicating impaired spatial memory. The study also found increased expression of Na⁺/H⁺ exchanger protein (NHE1) in reactive astrocytes, microglia, and oligodendrocytes across various brain regions, including the cortex, corpus callosum, and hippocampus. This upregulation of NHE1 was linked to increased oxidative stress, axonal damage, and gliosis following rmTBI.

1.9 Brief History of NMR

Nuclear magnetic resonance (NMR) was independently discovered in 1946 by Felix Bloch and Edward Purcell, who demonstrated that atomic nuclei with a magnetic moment can interact with electromagnetic radiation when exposed to a magnetic field. This discovery led to the development of NMR principles and mathematical models, which laid the foundation for the technique. Bloch and Purcell, along with their teams, showed that by applying a magnetic field and radiofrequency pulses, atomic nuclei could be manipulated and their emitted signals analyzed to gather information about molecular properties.

In the 1950s, Erwin Hahn furthered the technology by introducing continuous wave NMR, allowing measurement of nuclear relaxation times. The 1960s and 1970s saw Richard Ernst and colleagues introduce Fourier transform (FT) techniques, which greatly improved the speed, resolution, and

sensitivity of NMR spectroscopy. Alongside this, Ernst and Jean Jeener proposed two-dimensional NMR techniques, such as COSY and NOESY, which enabled detailed structural analysis by identifying atomic connections and spatial arrangements. In the early 1970s, Paul Lauterbur and Peter Mansfield pioneered the use of NMR for imaging. Lauterbur introduced the concept of using magnetic field gradients for spatial encoding, while Mansfield developed image reconstruction techniques, leading to the birth of magnetic resonance imaging (MRI), a crucial tool in medical diagnostics. Advancements in NMR technology, such as stronger magnets, advanced pulse sequences, and better data processing, have allowed NMR to be applied in diverse fields, including chemistry, biochemistry, materials science, and medicine.

1.9.1 Physical Basis of NMR

Nuclear Magnetic Resonance (NMR) relies on the behavior of atomic nuclei in a magnetic field. The process hinges on the interaction between nuclear spins and an external magnetic field, providing insights into the chemical and physical properties of molecules.

Nuclear Spin: Nuclei possess intrinsic angular momentum, known as spin, which is vital for NMR. Nuclei with odd numbers of protons or neutrons have a non-zero spin, making them suitable for NMR. Common nuclei studied include hydrogen (^1H), carbon-13 (^{13}C), and phosphorus-31 (^{31}P).

Magnetic Moment: The spinning nucleus generates a magnetic moment, which is a vector indicating the strength and direction of the magnetic field.

Zeeman Effect: When exposed to a magnetic field, the nuclear magnetic moment aligns either parallel (low energy) or antiparallel (high energy) to the field, resulting in a small energy gap between these two states, known as Zeeman splitting.

Resonance: A second magnetic field, a radiofrequency (RF) field, is applied to induce resonance, exciting the nuclear spins and transitioning them between energy states.

Larmor Frequency: The resonance frequency is determined by the Larmor equation, which links it to the external magnetic field and the gyromagnetic ratio of the nucleus.

Relaxation: After excitation, nuclear spins return to equilibrium via two processes: longitudinal relaxation (T_1) and transverse relaxation (T_2). T_1 represents the recovery of alignment with the magnetic field, while T_2 is the loss of transverse magnetization, leading to signal broadening.

NMR Signal: Once the spins relax, they emit a signal detectable by an NMR coil, which is processed to yield an NMR spectrum that provides molecular structure and dynamics information.

Net Magnetization

In magnetic resonance, the hydrogen nucleus (^1H) is commonly used to gather information in biological tissues due to its prevalent presence in the body. When placed in a magnetic field, the individual hydrogen protons align and precess at a frequency proportional to the field strength. The energy difference between protons aligned with or against the field is governed by the Zeeman interaction and described by the Boltzmann distribution.

The application of an RF pulse at the Larmor frequency can excite the protons, causing them to flip between energy states. The manipulation of these magnetic moments forms the basis of MR techniques.

Relaxation

Relaxation in NMR refers to the processes by which nuclear spins return to their equilibrium states after being disturbed by an RF pulse. These processes are:

Longitudinal Relaxation (T_1): Involves the return of spins to their original alignment with the magnetic field, typically via energy exchange with the surrounding environment.

Transverse Relaxation (T_2): Refers to the decay of the transverse magnetization, caused by interactions among spins and field inhomogeneities. Both T_1 and T_2 times affect the quality and resolution of NMR signals, and their characteristics depend on the molecular environment.

Free Induction Decay (FID)

The FID is the signal generated immediately after the application of an RF pulse, as nuclear spins precess and induce voltage in the NMR coil. The signal decays due to relaxation processes, and its decay is governed by T_2 . Fourier Transform (FT) is then applied to convert the FID from a time-domain signal into a frequency-domain spectrum, revealing valuable information about the chemical environment of the sample.

Chemical Shift (δ)

Chemical shifts arise due to variations in the magnetic field around a nucleus, primarily influenced by the surrounding electrons and other nuclei. Each proton experiences a unique local magnetic field, which shifts the resonance frequency. This shift is measured relative to a reference compound and is expressed in parts per million (ppm), enabling comparison across different NMR experiments.

Signal Intensity

The intensity of NMR signals correlates with the concentration of the observed nuclei. The area under a peak reflects the number of contributing nuclei, although relaxation effects can influence the measured intensity.

Spin-Spin Coupling (Multiplicity)

Spin-spin coupling refers to the splitting of NMR signals due to interactions between adjacent nuclei.

The number of peaks and their spacing depend on the number of neighboring nuclei. This coupling provides insights into molecular structure by revealing the number and type of nearby protons.

NMR Sequences for Biological Samples and In-Vivo MR Spectroscopy

Various pulse sequences are used in NMR for biological samples, with each chosen based on the type of data required:

Carr-Purcell-Meiboom-Gill (CPMG): This sequence is used to measure T_2 relaxation times, ideal for analyzing serum samples and detecting smaller metabolites.

ZGPR (Zero-Go-Pulse-Relaxation): A basic sequence often used to study water-solvent peaks in biological tissues.

Nuclear Overhauser Effect Spectroscopy (NOESY): This technique enhances the detection of metabolites by analyzing small changes in electron spin populations, commonly used for urine sample analysis.

1.10 NMR Metabolomics in TBI

Nuclear Magnetic Resonance (NMR) metabolomics has emerged as a valuable tool in traumatic brain injury (TBI) research, providing insights into the metabolic alterations associated with TBI.

NMR metabolomics enables the comprehensive analysis of metabolites present in biological samples, such as blood, cerebrospinal fluid, or brain tissue. By comparing the metabolic profiles of TBI patients with healthy individuals, researchers can identify specific metabolic changes associated with TBI. This approach helps in understanding the underlying metabolic pathways and dysregulations involved in TBI pathology. It allows the identification of potential biomarkers for TBI diagnosis, prognosis, and treatment monitoring. By examining the concentrations of specific metabolites or metabolic patterns,

researchers can discover biomarkers that reflect the severity of injury, predict patient outcomes, or monitor the effectiveness of therapeutic interventions. These biomarkers can aid in early detection, personalized treatment, and assessment of TBI progression. NMR metabolomics data can be used to investigate the perturbations in metabolic pathways following TBI. By integrating metabolomics data with other omics approaches, such as genomics and proteomics, researchers can gain a comprehensive understanding of the molecular mechanisms underlying TBI. This information helps in identifying key metabolic pathways involved in TBI pathophysiology and potential therapeutic targets. TBI often leads to secondary injury mechanisms, including neuroinflammation, oxidative stress, and altered energy metabolism. NMR metabolomics provides a means to assess these secondary injury processes by monitoring changes in metabolite levels associated with inflammation, oxidative damage, and energy metabolism. This knowledge can contribute to the development of targeted interventions to mitigate secondary damage and improve patient outcomes. Therapeutic Intervention and Monitoring: NMR metabolomics can be used to evaluate the effects of therapeutic interventions in TBI. By analyzing metabolic changes pre- and post-treatment, researchers can assess the efficacy of interventions and gain insights into the mechanisms of action. This information is valuable for optimizing treatment strategies and monitoring the metabolic response to therapy.

NMR metabolomics has the potential to enhance our understanding of TBI by providing valuable insights into the metabolic alterations associated with the injury. By identifying biomarkers, elucidating metabolic pathways, and assessing treatment responses, NMR metabolomics contributes to the development of personalized and targeted approaches for TBI diagnosis, prognosis, and therapeutic interventions.

Chapter 2

Effect of closed head injury in hippocampal metabolome of rats at acute time-point

2.1 Introduction:

Road accidents, falls, and assaults are the most common causes of blunt traumatic brain injury (TBI), which is highly prevalent in civilian populations. Survivors of TBI often experience long-term personality changes and impairments in cognitive and motor functions, highlighting the urgent need for novel pharmacological treatment options.

Research has shown that TBI involves multiple pathological mechanisms, including oxidative stress, excitotoxicity, membrane disruption, neuronal injury, energy failure, and mitochondrial dysfunction. Additionally, altered metabolite concentrations in the hippocampus have been linked to dysregulation of energy metabolism and excitatory neurotransmission. Metabolomics, which provides a comprehensive fingerprint of biochemical processes, is a valuable tool for studying complex diseases like TBI, offering insights into the underlying metabolic disruptions and potential therapeutic targets.

In an in vivo study Xu et al. (2011) observed lower levels of total creatine ratios of N-acetylaspartate, glutamate, myo-inositol, phosphocholine, glycerophosphocholine, and taurine on the ipsilateral side of the brain after 2 and 4 hours of injury in an MRS study on the CCI model of TBI. At 4 hours after TBI, there were significant decreases in myo-inositol and taurine, as well as minor reductions in NAA. Another study on the FPI model reported a decrease in glutamate, ascorbate, phosphocholine, glycerophosphocholine, and N-acetylaspartic acid (NAA) levels in the cortex and hippocampus after 1 h of injury (Viant et al, 2005). Singh et al, 2017 reported alterations in the concentration of hippocampal neurometabolites at 4hr and day1 after injury in mild and moderate TBI rats. They reported decreased level of BAA, lactate, succinate while an increase in the levels of acetate,

myo- inositol, taurine, GABA, glutamate. Importantly, TBI induced impairment in the neuronal metabolism is critical to the changing energy needs of the brain. A continuous energy flow, or adenosine triphosphate (ATP) production, is required to maintain basic neurological function and to also prevent the propagation of secondary injury cascades. Pandya et al. (2019) conducted a study on severe penetrating ballistic like brain injury model at acute timepoint and they reported decreased mitochondrial function and significant decrease in the level of total flavin adenine dinucleotide (FAD) and nicotinamide adenine dinucleotide (NAD) in frontal cortex and striatum of injured rats compared to sham after 24hrs of injury. Another study on moderate lateral fluid percussion injury reported significant decrease in cerebral blood flow at 6 and 24hrs after injury and decreased mitochondrial function after 6 hrs of injury (Jiang et al, 2000). Rana et al. (2020) found that altered metabolites in the hippocampus and frontal cortex are related to energy metabolism dysregulation and neurotransmitter metabolism at day 1, day 3, and day 7 of blast-induced traumatic injury using untargeted ¹H-NMR. An experiment on the controlled cortical impact (CCI) model of TBI using LCMS metabolomics reported alterations in arginine and proline metabolism as well as taurine and hypotaurine metabolism on day 3 post-injury in the rat hippocampus. Another study using LCMS reported alterations in the purine metabolism pathway in the cortical tissue of CCI rats at the sub-acute time point (Li et al.,2020).

Graded traumatic brain injury (TBI) in animal models is a research approach used to mimic the different levels of injury severity observed in human TBI cases. This technique allows researchers to study the effects of TBI across a spectrum of injury severities, ranging from mild to moderate to severe, and provides valuable insights into the underlying pathophysiological mechanisms and potential therapeutic interventions. The present study is designed to understand the pathophysiology of diffuse graded TBI on the metabolic homogeneity of the hippocampus at acute time point using ¹H-NMR metabolomics.

To generate diffuse injury in TBI, we used a modified version of Marmarou's weight drop paradigm and ¹H-NMR spectrometry to investigate metabolic changes in hippocampus regions following graded injury. Using ¹H-NMR metabolomics, this study may help to understand the metabolic alterations and pathophysiology of hippocampal tissues after 24hrs of closed head injury in the hippocampus.

2. 2 Material and method

2.2.1 Chemicals

Saline (0.9% NaCl), Ketamine (Psychotropics India Limited, India), and Xylazine (Indian Immunological Limited, India), 4% formalin, 4% paraformaldehyde fixative, Phosphatebuffered saline (pH 7.4), dH₂O, sodium chloride (NaCl), potassium dihydrogen phosphate (KH₂PO₄), acetonitrile (Sigma Aldrich, Germany), potassium chloride (KCl), trimethylsilylpropanoic acid (TSP) (Eurisotop, France), sodium hydrogen phosphate (Na₂HPO₄) (Merck, Germany) Thioflavin S, deuterium oxide (D₂O) (Sigma Aldrich, Germany), and liquid nitrogen were used.

2.2.2 Study model:

Total sixty-five (n=65) male Sprague-Dawley rats of age 10-12 weeks were acquired from the institutional experimental facility. The weight of animals ranges from 250-280 gm each. Prior to experiment the animals were housed in controlled conditions at least for a week at 12-hour day- night cycle, 22±2 °C temperature and 50± 10% of relative humidity with constant supply of food and water *ad libitum*. All the experiments in this study were done according to the ethical guidelines of the institutional animal ethical committee.

2.2.3 Injury Model:

Animals were randomly divided into injury groups (control – 5; each injury group- 10). To develop the injury models an in- house modified version of Marmarou's impact acceleration device was used. For injury a brass rod weighted 450gms and 15cm height with 2.5 cm diameter was freely dropped from different heights ranges 25cm to 150cm on the brain. Prior to injury the rats were anesthetized, and a stainless-steel button was placed on the sagittal suture to create a diffuse injury to the brain. Following injury, the animals were kept warm on a bed till they regained consciousness.

2.2.4 Probit analysis:

Injury severity was categorized on the basis of probit analysis, a statistical method widely used in radiation survival curves to estimate parameters that describe the response of a biological system to radiation exposure. This represents the relationship between the radiation dose and survival probability of the biological system. In present study, it represents the relationship between injury height and animal survival. Probit analysis helps to estimate the lethal dose (LD50), which represents the injury height that results in a 50% survival rate.

Based on the survival curve of the animals after injury, the injury heights were categorized as mild (25cm & 50cm), moderate (75cm), severe (100cm, 125cm & 150cm) and lethal (175cm & 200cm). The 175 cm and 200cm injury heights were excluded from the study because a statistically desired number of rats did not survive the injury from these two heights.




Figure 2.1: Survival curve following 24hrs of injury

2.2.5 Sample collection and processing:

After 24 hours post-injury, the animals were anesthetized, euthanized, and perfused with phosphate-buffered saline (PBS, pH 7.4) followed by 4% paraformaldehyde (PFA). After euthanasia, the brains were carefully extracted from the skull and dissected on ice. The hippocampus was isolated and immediately snap-frozen in liquid nitrogen before being stored at -80°C. Metabolites were extracted by mixing acetonitrile and water in a 1:1 (v/v) ratio. The tissues were homogenized using a handheld homogenizer (IKA T-10, Germany) at a low temperature to prevent metabolic degradation, with 5 mL of acetonitrile/water mixture per gram of frozen tissue. The homogenate was then centrifuged at 12,000 rpm at 4°C for 20 minutes. The supernatant was separated and stored at -80°C until lyophilization. After freeze-drying, the resulting powdered samples were stored at -20°C until further analysis by NMR spectroscopy. The whole brain was preserved in 4% formaldehyde for subsequent histological examination.

2.2.6 NMR data acquisition and processing:

The ¹H NMR spectra were recorded using a 600.33 MHz Bruker BioSpin spectrometer (Switzerland). The lyophilized samples were reconstituted in a mixture of 600µL D₂O and 0.5 mM trimethylsilylpropanoic acid (TSP). The solution was then centrifuged at 12,000 rpm at 4°C for 10-15 minutes. Following centrifugation, 580 µL of the supernatant was carefully transferred into 5 mm NMR tubes for spectral analysis. NMR spectra were acquired using a water-saturation NOESYPR1d pulse program at 298 K with the pulse sequence RD-90 -t- 90 tm- 90 -acq. A total of 64 scans, including four dummy scans, were performed with a spectral width of 9009 Hz, an acquisition time of 3.63 s, and a relaxation delay of 4 s. After acquisition, the raw data were processed using Bruker TOPSPIN 3.5 software. Phase and baseline corrections were applied manually, and the peaks were integrated. The

total sum of integrals for each spectrum was then utilized to normalize the peaks, with TSP serving as the reference for peak integration.

Histological staining

The fixed brain tissues were embedded in paraffin wax following processing. Following microtome the silane-coated silted were used to mount the 5 μ m thick coronal slices. The tissue sections underwent deparaffinization with xylene and were rehydrated through a graded ethanol series before being washed in water.

For staining, a 1% thioflavin S solution was prepared by mixing it with 100% alcohol in a 1:1 ratio. The slides were incubated with the thioflavin S solution in the dark for one hour. After staining, the sections were examined under a fluorescence microscope (LMI BM-Prime, London, England), and images were captured using a 20X objective lens.

2.3 Statistical analysis:

Metabolic alterations between groups were compared using one-way analysis of variance (ANOVA). For posthoc analysis, the LSD test was used. $p < 0.05$ was considered statistically significant (SPSS version 20).

2.4 Results:

The metabolic profiles of the hippocampus were examined to determine the blunt-induced modification in the brain after 24 hours of injury. A total of 18 metabolites were detected. These metabolites are involved in a wide range of metabolic processes, including energy metabolism, ketone body metabolism, osmoregulation, neurotransmission, and membrane integrity. (Figure2.2)




Figure 2.2: Representative NMR spectra of hippocampal tissues showing selected metabolites.

2.4.1. Metabolic changes in hippocampal tissues after graded injury.

In our samples, we observed changes in neurometabolites (GABA, glutamine, and glutamate). All these three neurometabolites were down-regulated in hippocampal tissues on 24hrs after injury at all injury heights. Although significant downregulation was observed in glutamate levels at 25cm, 50cm, 100cm, 125cm, and 150cm injury groups. Whereas glutamine levels were notably low in 25cm, 50cm, and 125cm injury height groups. While GABA levels were significantly downregulated at all the injury heights after injury except 150cm injury group.

The energy metabolites which were altered in our samples were lactate, succinate, and creatine. Lactate and creatine levels were significantly low in 25cm, 50cm and 25cm, 50cm and 100cm injury height groups respectively. Whereas a significant increase in succinate levels in 100cm and 150cm injury groups was observed at 24hrs post-injury.

The acetate levels were up-regulated at all injury heights at 24hrs after injury.

Taurine levels in hippocampal tissues were significantly down-regulated at 25cm, 50cm and 125cm injury heights after 1 day of injury. The levels of myoinositol was decreased at 25cm, 50cm and 125cm injury height groups significantly.

The tyrosine level was significantly up-regulated in the hippocampus in 25cm injury height group. The NAA (an amino acid derivative) levels in hippocampal tissues were significantly down-regulated at all injury heights except 125cm group after 24hrs of injury. BAA levels were significantly higher in 25cm, 50cm, 100cm and 150 cm injury groups after 24hrs of injury in compared to controls.

The choline level in 25cm, 50cm and 150cm injury groups were significantly lower after injury in compared to control rats.




Figure 2.3: Showing metabolic change in hippocampal tissue after 24hrs of injury.

Neurite plaques

Thioflavin S staining was performed to stain neurofibrillary tangles. We observed neurite plaques at all the injury heights except 25cm injury group rats. The number of neurite plaques was found to be in increasing order with injury severity. The maximum number of plaques was observed in 150cm injury height group after injury.




Figure 2.4 : Showing the presence of neurite plaques in hippocampus after 24hrs of injury.

2.5 Discussion:

This study explores the impact of various diffuse injury levels on animal survival and neural metabolism in hippocampal tissue at an acute time point using NMR metabolomics. After 24 hours post -injury, significant changes were observed in the levels of neurotransmitters, osmolytes, and energy metabolites in the hippocampus compared to control animals. Immunohistochemistry and thioflavin staining were also conducted across all injury severities. The data revealed a notable reduction in glutamate, GABA, glutamine, acetate, NAA, and BAA levels on day 1 post -injury, which may reflect disruptions in energy homeostasis. Given that the hippocampus consists primarily of glutamatergic neurons (approximately 90%), the observed decrease in glutamate could result from altered glutamate homeostasis following the injury (Briend et al., 2020; Piao et al., 2019). In normal brain physiology, the continuous transport of glutamine from astrocytes to neurons for glutamate and GABA production is essential for maintaining metabolic stability. (Albrecht et al, 2010). A

significant decrease in GABA level was observed on day 1 of injury in almost all the injury severity groups. Down-regulation in GABA levels suggest alteration in GABA production as an effect of injury. While significant decrease in the levels of glutamine in mildly injured groups may influenced by the disturbance in glutamate- glutamine cycle. The glutamate glutamine cycle is a critical process for the compartmentalization of glutamate and glutamine between neurons and glia, and during glutamatergic transmission, neurons release glutamate into the surrounding extracellular space (Petroff, 2007) (Sowers et al, 2021). Additionally, the level of NAA was significantly down regulated and acetate levels were notably high in hippocampal tissues after injury at almost all the injury heights. NAA acts as an acetate reservoir in the brain after injury to maintain acetyl-CoA levels (Moffet et al, 2013). This acetyl-CoA then enters the TCA cycle for energy production (Akram, 2014). Di Pietro et al. (2014) reported a decrease in whole brain tissue NAA levels in rats following mild and severe diffuse TBI using HPLC (Pietro et al, 2014). An NMR study on blast-induced repetitive mild TBI reported a decreased level of NAA in hippocampal tissues after 24 hours of injury (Kumari et al, 2023). This suggests NAA can participate in energy production with acetate as an energy substrate after TBI. Decrease in NAA level and increase in acetate levels in almost all the injury groups may due to high energy demand of the hippocampal tissues after injury.

AA is crucial for neurotransmitter synthesis and energy metabolism, and its levels decrease one day post-injury compared to controls (Sperringer et al., 2017). Similarly, our findings show a reduction in BAA levels in hippocampal tissue at 1 day post-injury (PI). This decline in BAA suggests that the tissue experiences a high energy demand shortly after injury. Taurine, a neuromodulator that regulates dopamine in the brain (Kontro et al., 1987), plays a role in the reward system. It is known to enhance dopamine release, and we observed lower taurine levels in both mild and severe injury groups compared to controls. Additionally, choline levels were reduced, particularly in the mildly injured rats. Choline is an essential phospholipid for cell membranes, necessary for synthesizing phosphatidylcholine and sphingomyelin, and for acetylcholine production, which is critical for cholinergic neurotransmission. A

loss of choline in the central nervous system (CNS) can impair acetylcholine release, potentially leading to deficits in spatial cognition and memory within the hippocampus (Zeisel et al., 2003; Zeisel, 1992). The observed decrease in choline following TBI suggests a disruption in neuronal membrane integrity, which could contribute to altered cholinergic neurotransmission postinjury. Following TBI, cells' energy demands to maintain vital cellular activities increase dramatically. This imbalance in metabolism and energy dysfunction can alter overall metabolic processes (Carpenter et al., 2015) and increase lactate production for energy (Patet et al., 2015; Van Hall et al., 2009). In our study, we found that lactate levels were downregulated in all mildly injured rats compared to controls. The reduction in lactate levels shortly after injury highlights its critical role in ATP production. Creatine, another essential energy metabolite, also plays a vital role in tissues with high energy requirements, such as the brain and muscles. As a key ATP reserve, creatine is crucial for energy storage and release. Our study observed a decrease in creatine levels in mildly injured rats compared to controls (Andres et al., 2008; Brosnan and Brosnan, 2007). Additionally, Andres et al. (2008) reported that in CNS diseases, creatine metabolism and the creatine kinase/phosphocreatine system are compromised, as the brain requires high amounts of ATP to maintain membrane potentials and signaling capacity. Finally, we observed significantly higher succinate levels in the hippocampus of severely injured rats. This increase may be linked to mitochondrial damage, which triggers the production of reactive oxygen species (ROS). Under aerobic conditions, intracellular succinate is rapidly metabolized into fumarate. Under anaerobic conditions, elevated levels of extracellular succinate bind to and activate the G protein-coupled succinate receptor GPR91 on stellate cells and macrophages in the liver, which triggers inflammation (Littlewood-Evans et al., 2016; Mills et al., 2021). Hypoxia also increases succinylation (Gibson et al., 2015; Chen et al., 2017). It is also reported that in metabolic function, succinate accumulation induces mitochondrial respiration and changes the metabolic profile (Ehinger et al., 2016; Ives et al., 2020; Zhang et al., 2020; Avram et al., 2021). Huang et al. (2022) reported accumulation of succinate in the neuronal stem cells of ischemic rats (Huang et al. 2022).

BAA works as a nitrogen doner in the glutamate synthesis, even around 25% of nitrogen bases accept nitrogen from leucine (Yudkoff, 1997). High level of BAA and low level of glutamate in injured hippocampal tissues may due to the interruption in glutamate synthesis. In addition to this thioflavin staining were done to observe the amyloid beta (A β) fibril formation after injury. We observed amyloid plaques at all the heights after injury. An ELISA study on TBI reported accumulation of A β fibrils in mice cortical tissues after 1,3 and 7 days of injury (Washington et al. 2014). Chen et al (2004) reported A β plaques in swine white matter after rotational head injury at all the timepoints (i.e. from day3 PI to 6 month PI). We also found our results in line with these studies.

2.6 Conclusion:

The current study uses NMR metabolomics to investigate the effects of a range of diffuse closed head injuries on hippocampus metabolism at an acute time point. TBI-induced metabolic changes are visible in changed levels of neurotransmitters, energy metabolites, and neuromodulators in hippocampus tissues. Furthermore, neurite plaques were discovered at all damage heights investigated, however the patterns of change varied depending on the injury category. This study gives important pathological insights into diffused closed head injury in people and contributes to our understanding of the TBI pathophysiology.

Chapter 3

Acute and sub-acute metabolic change in different brain regions induced by moderate blunt TBI

3.1 Introduction:

Brain injury can lead to significant metabolic changes in the body, both acutely and sub acutely. A brain injury disrupts the normal functioning of the brain, which can impact various metabolic processes throughout the body. (Bartnik et al., 2007) During the acute phase of a brain injury, there is a surge in metabolic activity as the body tries to repair and recover from the damage. This higher metabolic workload is mostly sustained by glucose and other nutrients obtained from the bloodstream. The interaction of disturbed intracellular bioenergetic pathways is expected to play a role in degenerative brain disorders. These metabolic alterations can cause brain edema, bleeding, and ischemia, which are all dynamic states that persist even after the initial damage. Furthermore, the severity of the brain injury can determine the intensity of the metabolic changes. In cases of mild brain concussion, there may be a brief increase in glucose metabolism in certain areas of the cerebral hemispheres and the brain stem. (Jalloh et al., 2014) In contrast, severe head injuries are associated with a consistent decrease in cerebral oxygen metabolic rate (Sunami et al., 1989). This decrease in cerebral oxygen metabolic rate can lead to further complications and impairments in brain function. Additionally, it is important to note that the pathophysiological mechanisms of brain injury are complex and can vary depending on the type of injury. (Vink et al., 1994)

Rana et al. (2020) found that altered metabolites in the hippocampus and frontal cortex are related to energy metabolism dysregulation and neurotransmitter metabolism at day 1, day 3, and day 7 of blast-induced traumatic injury using untargeted H1-NMR (Rana et al., 2020a).

An experiment on the controlled cortical impact (CCI) model of TBI using LCMS metabolomics reported alterations in arginine and proline metabolism as well as taurine and hypotaurine metabolism on day 3 post-injury in the rat hippocampus (Zheng, Zhou, Feng, et al., 2020). Another study using LCMS reported alterations in the purine metabolism pathway in the cortical tissue of CCI rats at the sub-acute time point (Zheng, Zhou, Li, et al., 2020). McGuire et al. (2019) reported metabolic changes in the hippocampus, thalamus, frontal cortex, striatum, and brainstem after the fluid percussion injury (FPI) model after 60 days of injury. It was found that levels of creatine, glutamine, glutamate, lactate, myo-inositol, and taurine in hippocampal tissues were altered. They also reported notable changes in the levels of creatine and glutamine in the thalamus (McGuire et al., 2019). A study on mice's brains after 2 and 7 days of excitotoxic neuronal injury found that distinct brain regions (cortex, hippocampus, striatum, midbrain, cerebellum, and brainstem) show variable vulnerability to injury and highlight the spatial role of injury-induced mechanisms (Jaeger et al., 2015). Another study on the FPI model reported a decrease in glutamate, ascorbate, phosphocholine, glycerophosphocholine, and N-acetylaspartic acid (NAA) levels in the cortex and hippocampus after 1 h of injury (Viant et al., 2005). Importantly, TBI-induced impairment in the neuronal metabolism is critical to the changing energy needs of the brain. A continuous energy flow, or adenosine triphosphate (ATP) production, is required to maintain basic neurological function and to also prevent the propagation of secondary injury cascades. A study on TBI patients using micro dialysis showed an elevated lactate/pyruvate ratio in white matter over the first 3–7 days after injury (Guilfoyle et al., 2021). Along with this, it has been reported that TBI leads to a prolonged glucose depression that can induce conditions like Alzheimer's disease as glucose deprivation promotes amyloidogenesis and tau pathology and affects synaptic function (Xu et al., 2021). Therefore, injury-induced metabolic alterations in the brain are important to understand to prevent long-term neurological deficits.

Because of the distinct tissue composition of each brain region, the metabolome of each region differs, leading to differential responses to injury. The majority of the neurons in the hippocampus are

pyramidal. Nakatomi et al. (2002) reported regeneration of hippocampal pyramidal neurons in ischemic rats(Nakatomi et al., 2002). The thalamus is made up of excitatory projection neurons and inhibitory interneurons, while the stratum is made up of medium-spiny neurons and interneurons. A study on Parkinson's disease reported altered medium spiny neurons (Deutch et al., 2007). A mouse CCI model of TBI study reported a notably lower proportion of inhibitory input neurons after 24 hours of injury (Frankowski et al., 2022). Another study on a mixed diffuse and focal injury reported a change in inhibitory neurons after 8 weeks of injury (Carron et al., 2020).

The current study is designed to understand the differential effect of diffuse brain injury on the metabolic homogeneity of the regions distant from the site of impact (hippocampus, thalamus, and striatum) at acute, early sub-acute and sub-acute time points using $^1\text{H-NMR}$ metabolomics.

For TBI, we used a modified version of Marmarou's weight drop model to induce diffuse injury and $^1\text{H-NMR}$ spectrometry to assess the metabolic change in tissues following injury. This study may help to understand the metabolic changes and pathophysiology of TBI in the hippocampus, thalamus, and striatum after day 1 and from the site of impact (hippocampus, thalamus, and striatum) at acute, early sub-acute and sub-acute time points using $^1\text{H-NMR}$ metabolomics. For TBI, we used a modified version of Marmarou's weight drop model to induce diffuse injury and $^1\text{H-NMR}$ spectrometry to assess the metabolic change in tissues following injury. This study may help to understand the metabolic changes and pathophysiology of TBI in the hippocampus, thalamus, and striatum after days 1, 3, and day 7 of injury. This study may help in better understanding the metabolic cascade of pathological events in different brain regions following trauma.

3.2 Material and method

3.2.1 Chemicals

Saline (0.9% NaCl), Ketamine (Psychotropics India Limited, India), and Xylazine (Indian Immunological Limited, India), 4% formalin, 4% paraformaldehyde fixative, Phosphatebuffered saline (pH 7.4), dH₂O, sodium chloride (NaCl), potassium chloride (KCl), potassium dihydrogen phosphate (KH₂PO₄), sodium hydrogen phosphate (Na₂HPO₄) (Merck, Germany), acetonitrile (Sigma Aldrich, Germany), trimethylsilylpropanoic acid (TSP) (Eurisotop, France), deuterium oxide (D₂O) (Sigma Aldrich, Germany), and liquid nitrogen were used.

3.2.2 Study model:

All experiments in this work were conducted in accordance with the Institutional Animal Ethical Committee's ethical guidelines (INM/IAEC/2019/09/Ext1). To evaluate the effect size, we determined the η^2 value for each metabolite in all three brain areas (supplementary table

1). We obtained twenty (n = 20) male Sprague-Dawley rats aged 10 to 12 weeks from the Institute's Experimental Animal Facility. These rats weighed an average of 250-280 g each. The animals were housed in a well-ventilated room with regulated circumstances, including a temperature of 22±2°C, relative humidity of 50±10%, and a 12-hour day-night cycle. The rats were constantly given adequate amounts of food and water. Rats were divided into two groups: sham controls (n=5) and rats with TBI (n=15). Out of 15 rats, brain tissue samples were collected at 24hr, 3day and 7 days (n=5 at each time point).

3.2.3 Injury Model:

The head injury animal model was developed using the in-house modified Marmarou's impact acceleration model (Singh et al., 2016). Rats were anaesthetized, and a stainless-steel button was placed on the skull. The button was placed between the bregma and lambda on the sagittal suture. The

stainless-steel button was placed to create a global injury to the brain. For injury, a brass rod of 450g was freely dropped from 75cm height. After injury, rats were placed on a warm bed until they regained consciousness.

3.2.4 Sample collection and processing:

At each time-point after injury, the animals were perfused with PBS followed by PFA. Following perfusion, the brains were excised from cranium and then hippocampus, thalamus and striatum were separated on ice and snap frozen in liquid nitrogen and then stored at -80⁰C. The acetonitrile extraction method was used for the extraction of metabolites. A handheld homogenizer (IKA T-10, Germany) was used to homogenize the tissue in a (1:1, v/v) acetonitrile/water mixture of 5 ml per 1 g of frozen tissue. The homogenized tissue mixture was centrifuged for 20 min at 12,000 rpm at 4 °C. The supernatant was collected and stored at -80⁰ C before lyophilization. The tissue samples were lyophilized following extraction and stored at -80⁰C till spectra acquisition. During the processing, some of the samples were lost. (Supplementary table 1).

3.2.5 NMR data acquisition and processing:

¹H NMR spectra were acquired utilizing 600.33 MHz (Bruker, BioSpin, Switzerland) spectrometer. The lyophilized examples were reconstituted in a combination of 600 µl D₂O and 0.5 mM trimethylsilylpropanoic acid (TSP). The mixture was then centrifuged at 12000 rpm at 40°C for 10-15 minutes. After centrifugation, 580 µl of the supernatant was separated and transferred into 5 mm NMR tubes for the acquisition of NMR spectra. A water saturation NOESYPR1d pulse program was used for the acquisition of NMR spectra at a temperature of 298 K, and with the pulse sequence RD-90 -t- 90 -tm- 90 -acq. Total number of scans (ns= 64), including four dummy scans, were performed, with the spectral width 9009 Hz, the acquisition time 3.63 s, and the relaxation delay 4 s. After spectra

acquisition, the Bruker TOPSPIN 3.5 software was used to process the raw NMR spectra. Manual changes were made to correct the phase and baseline correction and then peaks were integrated (Figure 2). TSP was used as reference for integration. After integration, the peaks were normalised by dividing the total sum of integrals for each spectrum.

3.3 Statistical analysis:

Metabolic alterations between groups were compared using one-way analysis of variance (ANOVA). For posthoc analysis, the Bonferroni test was used. $p < 0.05$ was considered statistically significant (SPSS version 20). Pathway analysis was done using MetaboAnalyst 5.0 software.

3.4 Results:

The metabolic profiles of hippocampus, thalamus, and striatum tissue were evaluated to assess the blunt-induced alteration in the brain after 1 day, 3 days, and 7 days of injury. A total of 18 metabolites were observed. These metabolites are involved in a variety of metabolic processes like energy metabolism, ketone body metabolism, osmoregulation, neurotransmission, and as associates in different metabolic pathways and membrane integrity.

One-way ANOVA

Neurometabolites in different regions were studied at each time-point. Temporal alterations were observed in the metabolic levels (Figure 3.1). In addition to metabolite alterations in each region with time, we also observed significantly altered metabolites between different regions at the baseline study.

3.4.1 Change in metabolic levels among different brain regions in control

At baseline, we observed increased levels of NAA, myo-inositol, creatine, glutamate, and lactate in the hippocampus compared to both the thalamus and striatum; however, these metabolites showed no change between the thalamus and striatum. Choline, alpha-KG, GABA, acetate, and branched amino acid levels were significantly lower in hippocampal tissue compared to both the striatum and thalamus. Significantly increased levels of aspartate and tyrosine were observed in the thalamus compared with both the hippocampus and the striatum. Taurine levels were highest in striatal tissue, followed by the hippocampus, and then the thalamus.

3.4.2 Temporal changes in metabolites within a region after injury

In our samples, we found changes in GABA and glutamate. Glutamate levels were downregulated in hippocampal tissues on day 3 after injury. TBI group rats had significantly lower GABA levels in thalamus at day 1 and in striatum at days 1 and 3, respectively.

The energy metabolites that were altered in our samples were lactate, succinate, and creatine. In the hippocampal region, significantly decreased levels of lactate and creatine were observed at day 3 PI. Whereas a significant increase in succinate levels was observed in the hippocampus at day 3 PI. No significant change was observed in the levels of these three metabolites in the thalamus and striatum at any time point after injury.

On day 1, the acetate levels in the thalamus region were significantly lower. The level of alpha KG showed a down-regulation at day 1 and day 3 PI in striatal tissues after injury.

Taurine levels in the striatum were significantly downregulated at days 1 and 3 PI. The levels of myo-inositol were significantly decreased at day 3 in the hippocampus, while the thalamus and striatum showed significantly increased myo-inositol levels at day 1.

The tyrosine level was significantly up regulated in the hippocampus at day 7 PI. The NAA (an amino acid derivative) levels in hippocampal tissues were downregulated at days 3 and day 7 PI.

Although, it reached statistical significance at day 3 PI. BAA showed no significant changes in the hippocampus at any time after injury. BAA levels were significantly lower in the thalamus and striatum on day 1 after injury compared to controls.

The choline level in the thalamus was significantly lower after injury at all the time points, while in the striatum it was significantly downregulated at day 1 and day 3 PI compared to control rats.




Figure 3.1: Line graph of all selected metabolites in hippocampus, thalamus and striatum at all three time-points. P- Value < 0.05, considered significant.

3.4.3 Pathway analysis:

Most of the observed metabolites participate in alanine, aspartate, and glutamate metabolism; phenylalanine, tyrosine, and tryptophan biosynthesis; glutamine and glutamate metabolism; and taurine and hypotaurine metabolism. Out of these four pathways, the alanine, aspartate, and glutamate metabolism pathways were found to be altered in all three regions, with the maximum impact value among all pathways (0.71). The alanine, aspartate, and glutamate metabolism pathways were found to

be altered in all three regions at different time points; they showed alteration in the hippocampus at days 3 and day 7 PI, in the thalamus at days 1 and 7 PI, and in the striatum at day 1 PI. Phenylalanine, tyrosine, and tryptophan biosynthesis were found to be altered in the hippocampus at days 3 and 7 PI, in the thalamus at day 1 PI, and in the striatum at day 3 PI. Taurine and hypo-taurine pathways were found altered only in the striatum at day 1 and day 3 PI.

Effected Pathway	Total Compound	Hits	Raw p	FDR	Impact	Region	Time post injury
Alanine, aspartate and glutamate metabolism	28	7	0.006777	0.062888	0.70754	Hippocampus	3 day
Phenylalanine, tyrosine and tryptophan biosynthesis	4	1	0.041105	0.064532	0.5	Hippocampus	3 day
D-Glutamine and D-glutamate metabolism	6	2	0.053962	0.06849	0.5	Hippocampus	3 day
Phenylalanine, tyrosine and tryptophan biosynthesis	4	1	0.0039051	0.023248	0.5	Hippocampus	7 day
Alanine, aspartate and glutamate metabolism	28	7	0.01271	0.039327	0.70754	Hippocampus	7 day
D-Glutamine and D-glutamate metabolism	6	2	0.031212	0.046818	0.5	Hippocampus	7 day
Phenylalanine, tyrosine and tryptophan biosynthesis	4	1	0.0006695	0.0033606	0.5	Thalamus	1 Day
D-Glutamine and D-glutamate metabolism	6	2	0.005319	0.010325	0.5	Thalamus	1 Day
Alanine, aspartate and glutamate metabolism	28	7	0.0074221	0.013607	0.70754	Thalamus	1 Day
Alanine, aspartate and glutamate metabolism	28	7	0.011791	0.059592	0.70754	Thalamus	7 Day
D-Glutamine and D-glutamate metabolism	6	2	0.016252	0.059592	0.5	Thalamus	7 Day
Taurine and hypotaurine metabolism	8	1	0.0001498	0.0022008	0.42857	Striatum	1 Day
D-Glutamine and D-glutamate metabolism	6	2	0.0023005	0.015183	0.5	Striatum	1 Day
Alanine, aspartate and glutamate metabolism	28	7	0.015456	0.046369	0.70754	Striatum	1 Day
Taurine and hypotaurine metabolism	8	1	6.76E-05	0.0019086	0.42857	Striatum	3 Day
Phenylalanine, tyrosine and tryptophan biosynthesis	4	1	0.034514	0.16191	0.5	Striatum	3 Day

Table1: List of significantly altered pathway following moderate injury.

3.5 Discussions:

The present study demonstrates the effect of moderate diffuse TBI on the neural metabolism of different brain regions at acute and sub-acute time points using NMR metabolomics. The effect of injury on brain regions was non-uniform. The levels of neurotransmitters and osmolytes were altered in all three regions, whereas the levels of choline were altered in the thalamus and striatum. In addition to neurotransmitters and osmolytes, energy metabolites were altered in the hippocampus. While acetone levels were changed only in the thalamus. The level of taurine was found to be altered only in striatal tissues. Alanine, aspartate, and glutamate metabolism; phenylalanine, tyrosine, and tryptophan biosynthesis; glutamine and glutamate metabolism; and taurine and hypotaurine metabolism are important pathways that change after injury.

Alterations in metabolic pathways are the result of collective metabolic changes rather than a single metabolic alteration. Several metabolites are involved in alanine, aspartate, and glutamate metabolism, as well as glutamine and glutamate metabolism. These metabolites function as a neurotransmitter or neuromodulator and as an energy metabolite. Alteration in these metabolic pathways is responsible for an imbalance between excitatory and inhibitory neurotransmitters and perturbs energy metabolism (Estrada Sánchez et al., 2008). Succinate, oxo-glutarate, pyruvate, and fumarate are metabolites that participate in the above metabolic pathways as well as the tricarboxylic acid cycle (TCA cycle) (Martínez-Reyes & Chandel, 2020). As a result, alterations in these metabolites may affect the TCA cycle. Phenylalanine is a precursor of tyrosine, and tyrosine participates in dopamine, epinephrine, and norepinephrine (Daubner et al., 2011; Fernstrom & Fernstrom, 2007). Perturbation in phenylalanine, tyrosine, and tryptophan biosynthesis may influence neurotransmitter production and ultimately cause cognitive decline (L.-N. Wang et al., 2022). 3-(4-Hydroxyphenyl) pyruvate is another intermediary metabolite that participates in phenylalanine, tyrosine, and tryptophan biosynthesis. 3-(4-

hydroxyphenyl) pyruvate changes into fumarate in the cytoplasm and enters mitochondria to participate in the TCA cycle (Bénit et al., 2014). Taurine and hypotaurine metabolism start with the conversion of l-cysteine to taurine, and then taurine transforms into taurocholate and 5-glutamyl taurine.

Metabolic alteration among brain regions in controls

Metabolic alterations were observed among brain regions in control animals. We found an increased level of myo-inositol, creatine, glutamate, NAA, and lactate in the hippocampus compared to the thalamus and striatum. While taurine level in hippocampus was found significantly lower than striatum and thalamus. Most of these metabolites are involved in energy generation, indicating that energy demand in the hippocampus is higher than in other regions of the brain. A study on the energy efficiency of different brain regions also reported that the hippocampus needs more energy for its function than the thalamus (Moujahid et al.,

2014). Metabolically, hippocampal tissues act differently compared to the other two regions.

This may be due to the unique tissue architecture and functions of different brain regions. Neurogenesis is the process of generating new neurons. The dentate gyrus of the hippocampus is one of the important sites for neurogenesis. Growing knowledge of neurogenesis suggests that neurometabolites, energy metabolites, neuromodulators, and osmolytes play important roles in this process. Glutamate, an excitatory neurotransmitter in mature neurons, can influence immature neural cell proliferation and differentiation (Schlett, 2006). GABAergic mechanisms control differentiation and the timing of synaptic integration in the sub-granular zone (Ge et al., 2007). Neurogenesis is known to be energetically demanding, and studies have revealed that lactate, rather than glucose, is favored as a metabolic substrate in neurons to support their activity when faced with high energy demands (Matsui et al., 2017). Lactate plays a crucial role in modulating molecular pathways associated with neurogenesis, such as angiogenesis, neuronal excitability, and plasticity. These pathways collectively contribute to the survival and development of newly formed neurons (Álvarez et al., 2014; Morland et

al., 2017). Non-neuronal cells in the neurogenic niche, such as astrocytes, endothelial cells, oligodendrocytes, and microglia, which direct the process of neurogenesis, are also influenced by lactate in their metabolic and signaling pathways. Additionally, NAA is one of the most concentrated metabolites in the brain; studies suggest that NAA works as an acetate reservoir in the brain, and the reduced level of NAA is considered parallel to low ATP levels (Vagnozzi et al., 2007). Creatine is an amino acid that is mostly found in muscle and the brain and helps in energy production. Leem et al. (2018) reported that creatine and exercise prevent a decrease in neurogenesis in mice's brains. GABA, aspartate, choline, acetate, and BAA were found to increase in thalamic tissues compared to the hippocampus, followed by the striatum (Leem et al., 2018). The thalamus contains a large number of gabergic neurons and interneurons (Arcelli et al., 1997; Jager et al., 2019). A study reported the highest gabaergic TCA cycle flux in the thalamus and hypothalamus (Tiwari et al., 2013). Acetate, aspartate, and branched-chain amino acids are used by the brain for energy metabolism (Andersen et al., 2019; Mishkovsky et al., 2012; Simon et al., 1967). Although cholinergic neurons are distributed to almost all parts of the brain, a major population is present in the striatum (Lim et al., 2014). In cholinergic neurons, acetylcholine is formed from choline and acetyl coenzyme A (acetyl coA) (Ulus et al., 1989). While the thalamus contains cholinergic axons and dense cholinergic innervation, which contain a high amount of acetylcholine, striatal cholinergic neuronal activity plays a crucial role in motor control, memory, social behavior, and behavioral flexibility. Striatum contains a relatively high amount of taurine (Ruotsalainen & Ahtee, 1996). Taurine is a neuromodulator, and it regulates dopamine in the brain (Kontro, 1987). Striatum contains a population of dopaminergic neurons. Dopamine is a part of the reward circuitry in the brain. Studies report that taurine increases the release of dopamine in the brain. We also observed a relatively high level of taurine in the control striatum.

Metabolic alteration in the hippocampus following TBI

TBI studies report that it induces neurogenesis in the brain. Neurogenesis requires a lot of energy; the brain prioritizes energy homeostasis maintenance (X. Wang et al., 2016). To meet the high energy demands of neurogenesis, neuronal stem cells switch their metabolism from glycolysis to oxidative phosphorylation (Maffezzini et al., 2020).

The significant decreased levels of glutamate, creatine, lactate, and NAA at day 3 after injury in the hippocampus of injured rats may be due to altered energy homeostasis. As the hippocampus contains 90% of glutamatergic neurons, a decrease in glutamate levels could be due to altered glutamate homeostasis after injury (Briend et al., 2020; Piao et al., 2019). Physiologically, transportation of glutamine from astrocytes to neurons for the production of glutamate, or GABA, takes place in continuation; this balance is crucial for metabolic stability inside the brain (Albrecht et al., 2010). Glutamate to alpha keto-gluterate conversion is a reversible reaction where it enters the TCA cycle, and succinate is an intermediate of the TCA cycle (McKenna, 2007). Thus, down-regulation of glutamate and up-regulation of succinate at day 3 PI may indicate the preference of the brain for energy production. A decrease in the level of glutamate and increased levels of succinate at day 3 PI in hippocampal tissues implies neuronal damage and a preference for energy production. At day 7 PI, both metabolites start to come across the control, which may be due to recovery after injury. In addition to this, the level of NAA was significantly down regulated in hippocampal tissues after 3 days of injury. NAA acts as an acetate reservoir in the brain after injury to maintain acetyl-CoA levels (Moffett et al., 2013). This acetyl-CoA then enters the TCA cycle for energy production (Akram, 2014). Di Pietro et al. (2014) reported a decrease in whole brain tissue NAA levels in rats following mild and severe diffuse TBI using HPLC (Di Pietro et al., 2014). A NMR study on blast-induced repetitive mild TBI reported a decreased level of NAA in hippocampal tissues after 24 hours of injury (Kumari et al., 2023). This suggests NAA can participate in energy production with acetate as an energy substrate after

TBI. A decrease in the level of NAA at day 3 PI may be due to tissue damage and high energy demand after injury. We observed an increase in the NAA at day 7 PI, which can be an indication of recovery after injury. Additionally, the decrease in lactate and creatine levels at day 3 PI may be due to neurogenesis and high energy demand, respectively. Also, myo-inositol levels were down-regulated in hippocampal tissues at 3 days PI. Myo-inositol is a known osmolyte, which maintains cell volume and fluid balance in the body. Barkai et al. (1978) reported decreased myo-inositol levels in affective disorders (Barkai et al., 1978). A study of major depressive disorder patients reported decreased myo-inositol levels in the hippocampus (Shirayama et al., 2017). Notably, depression is a common outcome of TBI (Fromm et al., 2004). The initial decrease and then increase in myo-inositol levels indicate the fluid imbalance and their recovery, respectively. Additionally, tyrosine levels were also found to increase after 7 days of injury. Tyrosine is important for neurotransmitter production, and the level of tyrosine directly affects its level in tissues (Fernstrom & Fernstrom, 2007). A HPLC analysis of the hippocampus of PTSD rats showed a decrease in the level of monoamines after 1 hour of stress, while an increase in monoamine levels was observed after 7 days of stress induction (Harvey et al., 2006).

Effect of TBI on the thalamus at acute and sub-acute time points

GABA, acetate, and BAA were down-regulated, while myo-inositol levels were up-regulated in the thalamus after 1 day of injury. The thalamus contains interneurons, which are inhibitory and release GABA (Kelsom & Lu, 2013). A decrease in GABA levels may be due to damage to GABAergic neurons in the thalamus (McGeer et al., 1983). Zhang et al. (2017) reported a decrease in GABA levels in the ventrobasal thalamus of chronic inflammatory pain rat models using HPLC (Zhang et al., 2017). A significant decrease in GABA level was observed on day 1 of PI. Down-regulation in GABA levels suggest GABAergic neuronal damage as an effect of injury, while almost partially normal levels of GABA at day 7 PI indicate improvement in neuronal damage. Acetate plays an important role in energy production and acetylcholine synthesis (Kwok & Collier, 1982). As an effect of TBI, the brain's energy

demands increase, and to fill this energy demand, acetate is transported into astrocytes for energy production (Moffett et al., 2013). Another fate of acetate after TBI is the formation of acetylcholine; as reported previously, the level of acetylcholine increases after brain injury (Scremin et al., 2006). We observed a significant decrease in acetate levels in the injury groups after 1 day of PI compared to control rats. BAA plays an important role in neurotransmitter synthesis and energy metabolism. The level of BAA is found to decrease after 1 day of injury compared to control (Jeter et al., 2013). Similarly, we also found a decreasing level of BAA in the thalamus at acute time (1 day PI). Decreasing levels of BAA and acetate after injury indicate the high energy demand of the thalamus at acute time point. In contrast, myo-inositol levels were found to increase in thalamic tissues on day 1 of PI. Choline levels were significantly down regulated in the thalamus after injury at all-time points compared to controls. Thalamus contains cholinergic axons and dense cholinergic innervation (Huerta-Ocampo et al., 2020). Reports suggest TBI can generate acute and chronic effects on the cholinergic system (Shin & Dixon, 2015). Previous studies have reported altered levels of acetylcholine in the brain following TBI. Studies have reported a decreasing trend in acetylcholine moving from acute to chronic time points (McGuire et al., 2018). Our results of a temporally decreased level of choline in thalamus tissue are in line with the previous observations. A significant decrease in myo-inositol levels at the acute time point and choline levels at all the time points in PI may be due to neuronal tissue damage and irregular cell volume.

Effect of TBI on the striatum at acute and sub-acute time points

In the striatum, GABA and alpha-ketoglutarate levels were found to be downregulated at days 1 and day 3 PI. The down regulation of GABA levels in the striatum may be affected by gabaergic neuronal damage, as the striatum contains 95% of gabaergic neurons in rodents (Tepper & Koós, 2016). Alpha-KG is an intermediate in the TCA cycle and provides energy for cell processes (Liu et al., 2018). A decrease in alpha-KG levels indicates altered energy metabolism in the striatum. A decreased level of

alpha-KG may influence the downregulation of GABA. Alpha-KG serves as the metabolic precursor of GABA and glutamate (Shank & Campbell, 1982). Decreased levels of GABA and alpha-KG at day 1 and day 3 PI show striatal neuronal damage and energy demand to recover the damage. BAA plays an important role in neurotransmitter synthesis and energy metabolism; it acts as a nitrogen source for glutamine production inside the brain (Jeter et al., 2013). Aquilani et al. (2005) reported that short-term supplementation of BAA can enhance the cognitive ability of severe TBI patients (Aquilani et al., 2005). A significant decrease in striatal BAAs was observed at day 1 PI. A decrease in BAA levels at day 1 PI and an almost normal level of it implies its role in energy metabolism and neurotransmitter synthesis. Taurine is an antioxidant and has neuroprotective properties (Oh et al., 2020). A study on neuronal damage rats reported that taurine treatment increases the survival of rats following damage. Taurine was also found to increase the survival of the rats and reduce striatal lesions after damage induced by 3-nitropropionic acid (3-NP) in the striatum (Rivas-Arancibia et al., 2001). Therefore, this suggests that decreased taurine levels at day 1 and day 3 PI in the striatum could progress into oxidative damage. The initial decrease (at day

1 and day 3 PI) in the level of taurine indicates striatal damage. While a partial recovery in taurine levels at day 7 PI may be due to the initiation of striatal damage repair, a decrease in choline levels at day 1 and day 3 PI was also observed in the striatum. Choline is an acetylcholine precursor, and acetylcholine synthesis takes place in cholinergic neurons. The concentration of choline regulates the concentration of acetylcholine in the striatum (Ikarashi et al., 2002). A study on the FPI model reported increased acetylcholine concentrations following the injury at early time points (4 hr and 12 hr) (Saija et al., 1988). The decrease in choline levels at day 1 and day 3 PI could be due to cholinergic neuron damage.

3.6 Conclusion:

Our study reports changes induced after diffuse moderate blunt injury at acute and sub-acute time points using NMR metabolomics in different brain regions. Moderate TBI induced metabolic alteration is evident in altered level of neurotransmitter, energy metabolites and neuromodulator in all brain regions at different timepoints. In addition to this, evidence of metabolic dysregulation was found in all brain regions studied, the patterns of change observed in each brain region were regionally different. These results suggest that TBI can cause widespread and persistent metabolic changes. This study provides useful pathological insights into closed head injury in humans and expands our knowledge of pathophysiology of TBI.

Metabolites	chemical shift	multiplicity	HMDB id
BAA	0.918 - 1.0693	m	-
Lactate	1.31	d	HMDB0000190
Alanine	1.47	d	HMDB0000161
GABA	1.88	m	HMDB0000112
Acetate	1.9	s	-
NAA	2.009	s	HMDB0000812
Glutamine	2.12	m	HMDB00641
Glutamate	2.34	m	HMDB00148
Succinate	2.39	s	HMDB0000254
Aspartate	2.82	dd	HMDB0000191
alphaKG	2.99	t	HMDB0000208
Creatine	3.025	s	HMDB000064
Taurine	3.41	t	HMDB0000251
Glycine	3.55	s	HMDB0000123
PC/Choline	3.58	s	HMDB00097
Myo inositol	4.05	t	HMDB0000211
Tyrosine	6.8	d	HMDB0000158

Supplementary Table 1: List of selected metabolites and their peak details.

Chapter 4

Acute metabolic alterations in the hippocampus are associated with decreased acetylation after blast induced TBI

4.1 Introduction

Blast explosion and consequent blast related concussion is linked to destructive alteration in the brain leading to behavioral and cognitive impairments in the affected individuals (Clark et al., 2018) Importantly, soldiers deployed in Iraq war diagnosed with blast concussion showed symptoms of post-traumatic stress disorder (PTSD) and depression (Hoge et al., 2015).

Explosive detonation causes concussive blast waves, which are related with cerebrovascular injury as well as disturbance of the blood-brain barrier. Low level blast exposure induces structural and functional changes in the brain vasculature including alterations in the arterial smooth muscle layer (Gama Sosa et al., 2019). Changes in the cerebral blood flow and damage associated with blast wave results in oxidative stress and dysfunctional cerebral metabolism in the exposed rats (Rana et al., 2020b). Mild blast concussion may have serious outcomes if exposed repetitively, as a single insult increases cerebral vulnerability such that rats re-exposed to blast waves show increased oxidative stress, vascular pathologies and inflammation (Kamnaksh et al., 2014). On the other hand, military soldiers who experience minor and temporary symptoms of blast TBI are redeployed to their services, where they may sustain repeated mild injuries from reexposure to blast waves.

A repeated mild injury further possess the individual at a risk of severe ongoing secondary injury mechanisms in the central nervous system (CNS) along with long term outcomes like neurodegenerative changes and chronic traumatic encephalopathy (Stern et al., 2011). At the molecular level it has been found that blast induced repetitive mild TBI is accompanied by a complex pathology involving changes in the cerebral metabolism with increased vascular and oxidative damage (Kamnaksh et al., 2014). Cerebro-cerebellar hypometabolism is found as a major contributing factor

leading to chronic post concussive symptom in Iraq combat veterans with repetitive blast-trauma (Peskind et al., 2011). Similarly, Prins et al. (2013) reported a decrease in cerebral glucose metabolism in both parietal cortex and hippocampus in a subpopulation of children and young adults who experienced repetitive mild TBI (Prins et al., 2013).

Crucially, changes in brain metabolism can affect cellular gene expression via the mechanism of epigenetic modifications. Histone modifying enzymes employ cellular metabolites to change histone modifications, which rewrites gene expression (Katada et al. 2012). By controlling the amounts of acetyl CoA through the tricarboxylic cycle (TCA cycle), the content of glucose in the cell has been shown to affect the activity of histone acetyltransferase (HATs). Similarly, the methylation pattern of the cell is affected by the concentration of the co-factor FAD/FADH2 in the cell as well as the levels of SAM (S-adenosylmethionine) and SAH (Sadenosylhomocysteine) (Weinhold 2006). Furthermore, following moderate controlled cortical impact (CCI), a reduction in histone methylation and H3 acetylation is seen, particularly in the hippocampus region (Gao et al. 2006).

A large number of preclinical investigations on traumatic brain injury (TBI) have used male test subjects. Increasing the number of sex-balanced studies may result in more reliable human clinical trials and a better knowledge of disease causes. This study was specifically designed to evaluate metabolic changes in women's health. It seeks to investigate how repeated blast concussions alter the metabolism of exposed female rats. Furthermore, the study investigates the relationship between metabolic and epigenetic changes caused by repeated blast injury.

4.2 Materials and method

4.2.1. Chemicals

Acetonitrile (Sigma Aldrich, Germany), deuterium oxide (D_2O) (Sigma Aldrich, Germany), formalin, ketamine (Psychotropics India Limited, India), mylar membrane, potassium chloride (KCl), potassium dihydrogen phosphate (KH_2PO_4), sodium chloride (NaCl), sodium hydrogen phosphate (Na_2HPO_4)

(Merck, Germany), trimethylsilylpropanoic acid (TSP) (Eurisotop, France) and xylazine (Indian immunologicals Limited, India).

4.2.2 Study Design

All experiments adhered to the ethical guidelines outlined by the Institutional Animal Ethical Committee (INM/IAEC/2019/09/Ext1). The study aimed to investigate differences in neurometabolite concentrations between sham and blast-exposed groups. The null hypothesis (H_0) posited no significant differences between these groups, while the alternative hypothesis (H_a) indicated otherwise. With an α level of 0.05 and 80% test power, a two-tailed test determined a minimum sample size of 5 animals per group. Twenty female Sprague Dawley rats (10–12 weeks old, weighing 250–280 g) were sourced from the Experimental Animal Facility of the institute. Animals were randomly assigned into groups of five and housed in a controlled environment (temperature $22 \pm 2^\circ\text{C}$, humidity $50 \pm 10\%$) with a 12-hour light/dark cycle, providing ad libitum access to food and water.

4.3 Blast Exposure

A compression-driven gas shock tube was employed to generate shock waves, based on the protocol described by Mishra et al. (2016). Rats were divided into four groups: sham, mild TBI, moderate TBI, and repetitive mild TBI (rmTBI), with each group comprising five animals. Animals were anesthetized with 3.5% isoflurane in zero air (19.9–21.9% O_2) and positioned horizontally on a platform within the shockwave tube. The mild and moderate TBI groups were subjected to single shock waves of 100 kPa and 190 kPa, respectively. The rmTBI group received three 100 kPa shock waves at intervals of 0 minutes, 30 minutes, and 24 hours. Sham animals underwent identical procedures without exposure to shock waves. At 24 hours postinjury, rats were perfused sequentially with phosphate-buffered saline (PBS) and 4% paraformaldehyde (PFA) to minimize blood contamination during tissue extraction. The

hippocampus was isolated from one hemisphere, snap-frozen in liquid nitrogen, and stored at -80°C for subsequent analyses. The other hemisphere was fixed in 4% formalin for histological examination.

4.3.1 Immunohistochemistry (IHC) Analysis

Fixed brain tissues were processed and embedded in paraffin wax. Sections of 5 μ m thickness were prepared using a microtome and mounted on silane-coated slides. Sections were deparaffinized in xylene and rehydrated through a graded ethanol-to-water series. After rinsing with PBS, tissues underwent antigen retrieval by incubation in heated sodium citrate (85°C) for one hour. To suppress endogenous peroxidase activity, sections were treated with hydrogen peroxide (H₂O₂) for 15 minutes, following the protocol of the Mouse and Rabbit Specific HRP/DAB (ABC) Detection IHC Kit (ab64264). After rinsing with PBS, sections were blocked with a protein-blocking buffer for 10 minutes and incubated overnight at 4°C with primary antibodies in a humidified chamber. Specific antibodies used included GFAP (MA5-12023, Thermo) for astrocytic activation and AH3 (MBS9126034, My Biosource) for histone acetylation quantification. Tissues were washed with PBS and incubated for 10 minutes with the secondary antibody, followed by a 10-minute treatment with streptavidin-peroxidase. Sections were then exposed to DAB (3,3'-diaminobenzidine) for visualization, counterstained with hematoxylin for 30 seconds, air-dried, and cover-slipped. Slides were examined under a microscope (LMI BM-Prime, London, England) at 20 \times magnification, and images were captured. GFAP-positive and AH3-positive cells were manually counted in the dentate gyrus (DG) region of the hippocampus for mild, moderate, and repetitive TBI groups. Cell counts were performed using ImageJ software (Bethesda, USA), with regions of interest (ROIs) selected in each DG area based on the methodology described by Gao et al. (2006). Positive cell counts were normalized per mm².

4.4 Sample Preparation for NMR and Spectra Acquisition :

Metabolites were extracted from tissue samples using the acetonitrile extraction method. Frozen tissue samples were homogenized in a 1:1 acetonitrile-water mixture (5 mL per 1 g of tissue) using a handheld

homogenizer (IKA T-10, Germany) under cold conditions to prevent metabolic degradation. The homogenized solution was centrifuged at 12,000 rpm at 4°C for 20 minutes. The resulting supernatant was collected and stored at –80°C until lyophilized. Lyophilized samples were then stored at –20°C for subsequent analysis.

Proton nuclear magnetic resonance (^1H NMR) spectra were acquired using a 600.33 MHz NMR spectrometer (Bruker, Biospin, Switzerland). For spectral acquisition, the lyophilized samples were reconstituted in 600 μL of a solution containing deuterium oxide (D_2O) and 0.5 mM trimethylsilylpropanoic acid (TSP). The mixture was centrifuged at 12,000 rpm at 4°C for 10–15 minutes, and 580 μL of the supernatant was transferred into 5-mm NMR tubes.

The NMR spectra were obtained using a water-saturation NOESYPR1d pulse program with the following parameters: pulse sequence (RD-90°-t-90°-tm-90°-acq), 64 scans (including 4 dummy scans), spectral width of 9009 Hz, acquisition time of 3.63 seconds, and relaxation delay of 4 seconds. All spectra were recorded at a temperature of 298 K.

The raw NMR spectra were processed using TOPSPIN 3.5 software (Bruker). Baseline and phase corrections were manually adjusted for each spectrum. Peak integration was performed manually, and peaks were identified by referencing the Human Metabolome Database (HMDB) and prior literature. TSP was used as the internal reference for peak integration. The integrated peak data were normalized to the total sum of integrals for each spectrum.

4.5 Concentration of metabolites

Concentration of metabolites in tissues were calculated using the following formula

[Equation 1 (Tyagi et al, 2011)]:

$$C(x) = C(TSP) \times \frac{N(TSP) \times I(X) \times V(s)}{N(x) \times I(TSP) \times W(s)} \quad \text{Equation 1}$$

Where,

$C(x)$ - concentration of metabolite,

$C(TSP)$ - Concentration of TSP,

I_x – Intensity of metabolite

N_x - Number of protons in integrated signal,

N_{TSP} -Number of protons in TSP,

$W(S)$ - Weight of wet tissues in grams,

$V(S)$ - Volume of sample in NMR tube.

The concentrations of all metabolites, excluding branched-chain amino acids (BAAs), are detailed in Supplementary Table 2. Quantification of BAA concentrations was not performed due to overlapping peaks of valine, leucine, and isoleucine. Instead, their relative intensities are provided in Supplementary Table 2.

4.6 Statistical Analysis

Metabolic and immunohistochemical differences between groups were evaluated using oneway analysis of variance (ANOVA) for comparison. Tukey's HSD test was conducted for posthoc analysis, with statistical significance set at $p < 0.05$. All statistical analyses were performed using SPSS software (version 20). To explore the relationships among metabolites in each injury group, Pearson's correlation coefficients were calculated for the 19 identified metabolites across all samples using MetaboAnalyst software.

4.7 Results

Metabolic profiles of hippocampal tissue were analyzed to identify blast-induced alterations 24 hours post-injury. Nineteen metabolites involved in various metabolic pathways were identified. These included metabolites associated with energy metabolism (lactate, succinate, creatine, alpha-ketoglutarate), ketone body metabolism (acetate), osmolytes (myo-inositol, taurine), neurotransmitter amino acids (glutamine, glutamate, aspartate, GABA), branchedchain amino acids (valine, leucine,

isoleucine), and other amino acids (alanine, glycine, tyrosine). Additionally, N-acetyl aspartate (NAA) and membrane-associated metabolites (phosphocholine/choline, phosphoethanolamine (PE), ethanolamine) were identified.

4.7.1 Neurometabolic alteration among the groups

Significant changes in the concentrations of N-acetyl aspartic acid (NAA), glutamate, acetate, phosphocholine/choline (PC/choline), phosphoethanolamine (PE), ethanolamine, and creatine were observed only in the repetitive TBI (rmTBI) group compared to the sham group 24 hours post-injury. Acetate levels were significantly increased, while NAA, glutamate, PC/choline, PE, ethanolamine, and creatine were notably decreased in the rmTBI group. Additionally, lactate, alanine, and branched-chain amino acids (BAAs) displayed significant changes across all TBI groups when compared to the sham.

Lactate levels were elevated, whereas alanine and BAA levels were reduced in all injury groups.




Figure 4.1: Graph showing the significantly changed hippocampal tissue metabolites after brain injury.

4.7.2 Correlation among the neurometabolites

To explore the relationships among altered metabolites post-injury, a correlation heatmap was generated using MetaboAnalyst software. NAA was positively correlated with glutamate, creatine, ethanolamine, PE, and PC/choline, while showing a negative correlation with acetate and lactate concentrations.




Figure 4.2: Heat diagram of correlation analysis between metabolites.

4.7.3 Histological changes among the groups

Astrocytic activation following TBI was assessed using GFAP immunostaining. At 24 hours post-injury, the number of astrocytes in the hippocampus was significantly higher in the mild and moderate injury groups compared to the sham group (Figure 4.3). Although the rmTBI group showed increased astrocyte proliferation in the hippocampus, the change was not statistically significant.

To evaluate TBI-induced epigenetic changes, AH3-positive cells were examined. One-way ANOVA revealed a significant reduction in AH3-positive cells in the rmTBI group compared to the sham and mild TBI groups (Figure 4.4). No significant changes were observed in other groups.




Figure 4.3: GFAP positive cells in the dentate gyrus region of hippocampus of different injury groups and bar graph of GFAP positive cells at 24hrs PI.




Figure 4.4 : AH3 positive cells in the dentate gyrus region of hippocampus of different injury groups and bar graph of AH3 positive cells at 24hrs PI.

4.8 Discussion

Mild traumatic brain injury (mTBI) has garnered increasing attention in both clinical and preclinical research, yielding extensive data on post-injury metabolic changes in the brain (Banoei et al., 2018). mTBI is particularly prevalent among military personnel exposed to repeated blasts during training, combat, or field trials (Hoge et al., 2015). Mild blasts are known to impair cognitive, metabolic, and behavioral functions (Arciniegas et al., 2002; Subramaniam et al., 2015; Rana et al., 2020). However, the timeline for recovery and the benchmarks for evaluating these recoveries remain under discussion. This study assessed the metabolic changes in hippocampal tissue following different severities of injury (mild, moderate, and repeated mild blast exposures) at 24 hours post-injury. Significant alterations in the levels of NAA, acetate, PC/choline, PE, ethanolamine, glutamate, and creatine were observed exclusively in rmTBI rats compared to sham. These changes suggest that blast-induced metabolic disruptions may impact energy metabolism and epigenetic processes.

Histologically, the study found reduced acetylated cells in the hippocampus of rmTBI rats compared to sham animals. Histone acetylation is a crucial epigenetic mechanism, and the enzyme AceCS1 plays a vital role in this process. Previous studies indicate that acetate serves as an energy source under hypoxic conditions (Vazquez et al., 2016) and may trigger epigenetic changes (Gao et al., 2006). Consistent with earlier findings, this study observed increased acetate levels and reduced histone acetylation post-injury, suggesting acetate may be rerouted to energy production rather than epigenetic modification.

NAA's role in generating acetate through deacetylation by ASPA (aminoacylase II) aligns with the observed reduction in NAA levels post-rmTBI (D'Adamo Jr et al., 1972). Furthermore, decreased glutamate levels alongside reduced NAA concentration indicate impaired neurotransmission, consistent with findings from studies on Alzheimer's disease (Lin et al., 2003; Dedeoglu et al., 2004).

In astrocytes, the glutamate-glutamine cycle supports neurotransmission. This study reported decreased alanine and glutamate concentrations after rmTBI, indicating potential disruptions in this cycle. Increased GFAP-positive cells after mild and moderate injuries further underscore metabolic stress

following rmTBI. Additionally, reduced PC/choline levels in rmTBI rats point to compromised membrane integrity, potentially affecting cholinergic neurotransmission (Zeisel et al., 2003).

Lactate, a critical energy substrate, was elevated across all TBI groups, reflecting increased energy demand and glycolytic activity post-injury (Dienel, 2014). Creatine, essential for ATP production, was decreased in rmTBI rats, indicating disrupted energy metabolism (Andres et al., 2008). Similarly, reduced PE and ethanolamine levels suggest their role in energy production may be impaired following rmTBI (Karagezyan et al., 1975).

The study also observed reductions in BAA concentrations across all injury groups. BAAs are vital for energy metabolism and neurotransmitter synthesis, and their depletion post-TBI underscores potential metabolic stress (Caplan et al., 2018).

The metabolic alterations specific to the rmTBI group highlight the vulnerability of the brain to secondary injury mechanisms after repeated trauma. These changes reflect impaired mitochondrial function, oxidative stress, and metabolic shifts in energy production, which could lead to long-term neurological deficits (Roberto et al., 2008).

4.9 Conclusion

This study demonstrates that blast-induced TBI leads to significant metabolic changes at acute time points, particularly in the rmTBI group. Altered levels of NAA, glutamate, acetate, creatine, PC/choline, PE, and ethanolamine suggest a shift towards energy production at the expense of other cellular functions. Reduced acetylation in hippocampal cells post-rmTBI indicates an interplay between metabolic and epigenetic mechanisms. The findings also reveal increased astrocyte proliferation in mild and moderate injuries, with a modest increase in rmTBI. These results emphasize the need for further studies to explore the relationship between metabolism and epigenetic regulation in TBI recovery.

Supplementary table 1: Concentration of Metabolites (x10³ mmol/mg) in Sham and Injury groups

GROUP	tyrosin e	ml	Glycin e	Cho	Taurin e	PE	Ethano l-amine	Creatin e	alphaKG	Aspartat e	Succinat e	Glutamat e	GABA	Glutamin e		acetat e	Alanin e	Lactat e	BAA*
Sham1	5.433	405.637	0.165	1.172	1.156	3.624	0.320	3.979	2.738	1.293	0.090	5.292	2.408	5.889	3.171	0.662	0.339	6.580	14.864
Sham2	4.338	338.339	0.255	1.406	2.811	4.216	0.535	5.600	1.908	0.889	0.117	7.816	1.830	8.007	4.288	0.432	1.690	0.746	101.654
Sham3	3.470	336.544	0.297	1.600	3.169	4.764	0.638	6.016	2.052	1.285	0.095	7.722	1.466	6.791	5.680	0.322	1.426	0.604	75.295
Sham4	3.113	386.256	0.273	1.418	2.322	3.632	0.638	4.555	0.748	1.098	0.096	6.369	1.672	5.879	4.634	0.287	1.337	0.590	93.387
Sham5	2.393	352.694	0.280	1.128	2.276	3.094	0.408	4.550	1.381	1.061	0.096	6.480	1.334	6.625	3.374	0.360	1.320	0.645	93.602
m TBI1	14.936	303.200	0.202	1.094	2.566	3.188	0.503	4.454	1.889	1.007	0.160	5.171	2.137	6.180	2.889	0.706	0.789	7.132	16.958
m TBI2	-1.250	382.281	0.212	1.205	2.395	3.820	0.559	5.068	1.862	0.936	0.148	6.445	1.566	6.596	3.932	0.489	0.577	7.540	15.400
m TBI3	7.931	413.363	0.169	1.012	2.063	2.911	0.421	4.342	1.393	0.941	0.121	6.040	1.401	5.947	3.174	0.440	0.556	6.623	13.406
m TBI4	7.150	429.503	0.227	0.782	2.006	2.711	0.344	3.110	1.605	0.873	0.150	4.266	1.631	4.901	2.254	0.666	0.536	5.265	21.051
m TBI5	4.078	304.457	0.220	1.561	1.540	4.828	0.427	5.301	3.647	1.723	0.120	7.051	3.209	7.846	4.225	0.882	0.452	8.767	14.864
rm TBI1	1.230	430.659	0.311	1.211	2.171	3.768	0.318	5.010	1.633	0.714	0.085	5.932	1.562	6.766	3.719	1.320	0.599	9.163	9.552
rm TBI2	0.708	470.549	0.247	0.850	0.809	2.576	0.357	2.717	1.587	0.986	0.082	3.110	1.265	3.660	2.394	1.258	0.294	5.868	10.429
rm TBI3	0.702	321.872	0.266	0.680	1.242	1.689	0.332	2.471	1.060	0.685	0.039	3.636	1.046	3.970	1.878	1.031	0.371	4.334	9.488
rm TBI4	1.854	320.959	0.206	0.915	2.179	2.910	0.431	3.329	2.087	1.143	0.149	4.158	2.005	5.627	2.103	2.130	0.749	7.993	24.026
rm TBI5	12.403	289.971	0.241	0.977	2.091	3.340	0.409	3.686	1.805	1.068	0.144	5.013	1.926	6.146	2.451	1.819	0.675	7.644	20.741
mo TBI1	1.132	395.424	0.306	1.006	2.125	3.453	0.503	4.112	1.491	0.989	0.109	5.387	1.262	5.665	3.245	0.102	0.455	6.344	11.851
mo TBI2	-9.133	452.309	0.362	1.420	2.641	3.624	0.496	5.565	2.286	1.064	0.060	7.078	1.694	6.977	4.888	0.374	0.631	10.402	7.783
mo TBI3	-1.091	378.924	0.248	0.931	1.091	2.360	0.248	3.112	1.191	0.659	0.053	3.440	1.170	3.417	2.940	0.260	0.237	6.117	3.511
mo TBI4	2.451	356.561	0.318	1.343	2.658	3.852	0.405	5.184	1.923	1.009	0.107	7.010	1.692	7.221	4.128	0.374	0.698	9.588	10.391
mo TBI5	6.583	382.346	0.318	1.061	2.172	2.977	0.367	4.041	1.968	0.808	0.103	5.767	1.690	6.426	3.493	0.447	0.636	7.464	14.520

* Relative Intensity

Chapter 5

Closed head injury induced change in rat metabolome and gut microbiota

5.1 Introduction

Repetitive mild traumatic brain injury (rm TBI) is a condition in which a person sustains multiple mild traumatic brain injuries over time. Athletes participating in contact sports such as football, hockey, or soccer, and persons involved in activities involving a high risk of falls or head impacts are all at risk of repetitive mild TBI. Repeated mild TBIs can have a cumulative effect, which means that each consecutive damage may have a bigger influence on the brain's functioning and healing. rmTBI studies reported chronic metabolic, microstructural and pathological alterations along with cognitive decline in rodents (Boska et al; 2013). Vike et al, 2022 reported metabolic alterations in collegiate football athletes before and after sports season and they found alteration in xanthine metabolism, fatty acid metabolism, medium chain fatty acid, primary bile acid metabolism and glycolysis, gluconeogenesis, and pyruvate metabolism pathways. Another study on CCI mice models reported disrupted axonal transport and accumulation of APP on corpus callosum and prefrontal cortex at 48h, 1w, 4w and 10w post rmTBI. Same study highlights an increase in APP/beta- actine levels using western blotting at 8h, 48h and 1w after injury. Ravdin et al, 2003 examine the cognitive change for a month in professional boxers and found significant cognitive decline in them. Along with metabolic changes traumatic brain injury has been found to have a notable impact on the composition and functioning of the gut microbiome. More than 10^{13} - 10^{14} gut microorganisms from over 1000 different known bacterial species exist in the human gut, which outnumber the somatic cells in the human body, and the bacterial genome is 100-fold greater than those in the human genome. The gut microbiota has a close relationship with the central nervous system (CNS). Studies have demonstrated that traumatic brain injury can alter the diversity and abundance of gut microbial species, leading to dysbiosis and the development of virulent phylogenetic changes within the intestinal commensal microbiota. This disruption of the gut microbiome following traumatic brain injury is thought to be a result of the bidirectional

communication between the brain and the gut. This communication is mediated by various factors, including immune modulatory metabolites, gut peptides, neurotransmitters, and cytokines. Furthermore, it has been shown that the extent of the brain injury, as indicated by the volume of the lesion seen on

MRI, is significantly correlated with the severity of the changes in the gut microbiome. Additionally, research suggests that damage to distant organs, such as burn injury or ischemia reperfusion, can also lead to alterations in the gut microbiome (Kosakamoto et al., 2020).

Treangen et al. (2018) reported that traumatic brain injury (TBI), an acute brain injury, can lead to significant changes in the gut microbiome. Specifically, they observed a decrease in the abundance of *Lactobacillus gasseri*, *Ruminococcus flavefaciens*, and *Eubacterium ventriosum* and an increase in *Eubacterium sulci* and *Marvinbryantia formatexigens* 24 hours post-TBI. Additionally, the study demonstrated that gut microbiome dysbiosis induced by TBI persisted for up to 7 days and was correlated with lesion volume. Similarly, cerebral ischemic stroke, another form of acute brain injury, has been shown to induce gut microbiota dysbiosis, which is associated with injury severity and outcomes. These alterations in the gut microbiome not only worsen the pathological outcomes of TBI but also contribute to sterile inflammation. The gut microbiome—a complex community of microorganisms residing in the gastrointestinal tract—plays a critical role in regulating human health and disease. It has been increasingly recognized that the gut microbiome can impact brain function and normal brain growth (Hassani et al., 2022). Emerging evidence suggests that the gut microbiome may have an impact on neurological disorders such as Alzheimer's disease, Parkinson's disease, and traumatic brain injury. Specifically, studies have shown that the gut microbiome is involved in the regulation of pain perception and can contribute to the occurrence of neurologic disorders such as anxiety and autism. Moreover, alterations in the gut microbiota have been linked to inflammatory cascades and increased risk of developing post-traumatic stress disorder following traumatic brain injury (Li et al., 2022).

Research has demonstrated that there is a direct connection between gastrointestinal function and the brain, as seen in cases of traumatic brain injury (Ye et al., 2021).

The current study aimed to investigate TBI-induced alterations over time, from acute to chronic, following mild and rmTBI. The study used ¹H-NMR metabolomics to examine metabolic changes in serum. Fecal 16S rRNA sequencing were used to investigate the potential processes that alter following injury the gut microbiome.

5.2 Material and method

5.2.1 Chemicals

Acetonitrile (Sigma Aldrich, Germany), deuterium oxide (D₂O) (Sigma Aldrich, Germany), formalin, ketamine (Psychotropics India Limited, India), mylar membrane, potassium chloride (KCl), potassium dihydrogen phosphate (KH₂PO₄), sodium chloride (NaCl), sodium hydrogen phosphate (Na₂HPO₄) (Merck, Germany), trimethylsilylpropanoic acid (TSP) (Eurisotop, France) and xylazine (Indian immunologicals Limited, India).

5.2.2 Animal model

Thirty female SD rats weighing 200 to 250 g were randomly assigned to one of three groups: Control, Mild TBI, and rmTBI. For mild and rmTBI, rats were placed on a foam bed and subjected to a free-falling brass rod from a height of 25 cm, above the sagittal halfway of the rat brain. Rm TBI rats were subjected to three impacts at 24-hour intervals.

After the acclimatization of rats for seven days, the injury was induced, and all the experimental protocols were approved by the institutional ethical committee. The rats were kept at a temperature of 23±1 °C with a 12-h light/dark cycle and provided with a standard pellet diet (M/S Golden Feeds, Meharuli, Delhi, India) and water *ab libitum*.

5.2.3 Sample Collection and preparation

Serum

At 4hrs hours, 5day and 30day PI, blood (about 1 ml) was obtained from each animal's retroorbital vein in serum vacutainers, left to clot for 30 minutes at room temperature, and then centrifuged at 2500g for 15 minutes. The serum-containing supernatant was collected and kept at 80 °C for subsequent analysis.

Feces

The rats were kept in metabolic cages as they gained consciousness for the collection of urine and fecal pellets. The samples were collected at 30day and kept at -80°C for further analysis.

5.3 NMR spectra acquisition

As a reference, serum sample (450 μ l) was transferred to a 5 mm NMR tube containing 1 mM TSP in a closed capillary.

The NMR spectra were acquired at 600MHz, Bruker spectrometer (Bruker, Germany) at 300 K. For serum NMR acquisition, Water suppressed Carr-Purcell-Meiboom-Gill (CPMG) spin echo pulse sequence [RD-90°-(τ -180°- τ) n acquire] with a total spin echo (2n τ) of 200 ms was used to attenuate broad signals from proteins and lipoproteins. RD (relaxation delay) was 2.5 s, [RF (radio frequency) pulse], [τ (spin echo delay)], [n (number of loops)]. Typically, 128 scans were collected into 32 K data points over a spectral width of 12,820.51 Hz with an acquisition time of 9.29 s.

Data Processing

The serum spectra underwent a correction process that involved adjusting spectral phase, baseline, and chemical shift referencing for each individual spectrum. This correction procedure was carried out using TOPSPIN 3.7, a software developed by Bruker in Germany. Subsequently, the spectral region spanning from 0.5 to 9.5 ppm was quantified, and each specific region was normalized based on the total integral of the spectrum. The identification of metabolites was accomplished by referencing the human metabolome database (HMDB) and relevant literature.

Changes in the concentration of metabolites were compared among the control group, the mild TBI group, and the rmTBI group using a one-way analysis of variance (ANOVA). For each serum sample (450 μ l), it was transferred to a 5 mm NMR tube with a coaxial tube containing 1 mM TSP (Trimethylsilyl-2,2,3,3-tetra-deuteropropionic acid) as a reference compound within a closed capillary system.

5.4 Fecal 16S rRNA sequencing

DNA Isolation and QC

Genomic DNA of all the samples was isolated using DNeasy Power Soil Pro Kit (Qiagen). DNA purity was checked using NanoDrop 2000 (Thermofisher Scientific, Massachusetts, USA) and the integrity of DNA was evaluated on 1% agarose (Lonza, Belgium) gel. *16S rDNA V3-V4 metagenomics Amplification*. Total volume of 12.5 ng DNA from each sample was subjected to 16S V3-V4 Polymerase chain reaction (PCR) with the primers described below. The PCR products were then loaded on 1.5% Agarose gel along with a 100 bp ladder. Illumina adapter overhang nucleotide sequences were added to the gene-specific sequences (gene-specific sequences target the 16S V3 and V4 region).

The full-length primer sequences, using standard IUPAC nucleotide nomenclature (N=A/T/G/C, W=A/T, H=A/C/T, V=A/C/G), targeting the V3-V4 region:

16S	Amplicon	PCR	Forward	Primer:
			5' TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGNGGCWGCAG 3'	
16S	Amplicon	PCR	Reverse	Primer:
			5' GTCTCGTGGCTGGAGATGTGTATAAGAGACAGGACTACHVGGTATCTAATCC 3'	

After confirming the amplification on gels, 5 μ l of PCR reaction were pooled for each group, control, mild and repetitive mild. The pooled PCR products were bead purified, and eluted and 5 μ l of eluate was subjected to another round of PCR with dual indices and adapters to generate the respective libraries as recommended by Illumina. To determine the size range of the fragments and the average library size the cleaned libraries were quantitated on Qubit Fluorometer and appropriate dilutions loaded on a D1000 screen tape.

Data Analysis

All the analysis was carried out in R version 4.2.2. The package dada2 was used to read QC, read filtering, denoising, read merging, chimera removal and taxonomy assignment using the silva database. Mean quality score of more than 30 was considered of high quality. Multiple sequence alignment of representative sequences was done using the R package DECIPHER. The bioinformatics workflow included demultiplexing of the raw sequences which were then denoised to get the representative sequences for taxonomic classification and sequence alignment for phylogenetic studies. This data was then used for the assessment of differential abundance and diversity analysis. EdgeR, an R package that uses negative binomial distribution was used for comparative analysis. The comparisons were made based on the abundance of OTU between different sample groups.

5.5 Statistical Analysis

All values are expressed as mean \pm standard deviation (SD). The data were analyzed in IBM SPSS statistical software version 20 (IBM Corp., Armonk, N.Y., USA) using one-way analysis of variance (ANOVA) followed by post-hoc Tukey's LSD for comparison between groups. In addition, the relative intensity of metabolites was also analyzed using Metaboanalyst 5.0. Fecal sequencing data analysis was performed using R software. Values with $p < 0.05$ were considered significant.

5.6 Results:

Fecal Microbiome changes using 16S rRNA sequencing post blunt impact TBI

The gut microbiome composition was observed using 16S rRNA sequencing which showed changes in both mild and repetitive mild injured rats. The α diversity was expressed as Chao, Shannon, Simpson and Fisher indexes. The β diversity was evaluated using ordination analysis. The α - diversity showed an increase in richness within the mild group while rmTBI group showed decrease level of richness. At the phylum level, the Firmicutes and Bacteroidetes were decreased in rmTBI group. While phylum firmicutes was downregulated in mild TBI rats compared to sham. Mild injured rats showed increased Proteobacteria abundance.

The genus abundance showed decrease in *Agathobacter*, *Faecalibacterium*, *Holdemanella*, *Lactobacillus*, *Oribacterium*, *Rikenellaceae-RC9*, *Ruminococcaceae_UCG-008*, *Ruminococcaceae_UCG-014* and *Subdoligranulum* in mild and rmTBI groups. While *Anaerostipes*, *Fusicatenibacter*, *Marvinbryantia*, *Parabacteroides*, *Rambousta* and *Treponema* showed increase in mild and rmTBI rats compared to controls. *Alloprevotella*, *Bacteroides*, *Blacutia*, *Candidatus_Saccharimonas*, *Dorea*, *Prevotella_9*, *Prevotellaceae_NK3B31* group, *Pygmaeobacter*, *Ruminococcaceae_UCG-002*, *Ruminococcus_2*, *Solobacterium* and *Succinivibrio* showed increased abundance after mild

injury only. On the other hand, *Lachnospiraceae_ND3007_group*, *Lachnospiraceae_NK3A20_group*, *Lachnospiraceae_UCG_002*, *Oscillibacter*, *Sellimonas* and *Solobacterium* showed increased abundance only in rmTBI rats compared to control. The EdgeR analysis showed significant alterations in the genus *Blautia*, *Lachnospiraceae_ND3007*, *Lachnospiraceae_UCG_002*, *Lactobacillus*, *Ruminococcaceae_UCG-014* and *Sellimonas* from the phylum Firmicutes. From the Bacteroidetes phylum, genus *Alloprevotella*, *Parabacteroides*, *Prevotella_9*, *Pygmaiovacter* and *Rikenellaceae-RC9* showed significant change. *Alloprevotella*, *Parabacteroides*, *Prevotella_9*, *Pygmaiovacter* and *Candidatus_Saccharimonas* abundance was high in mild TBI rats, whereas *Lachnospiraceae_NK3A20_group*, *Lachnospiraceae_UCG_002* and *Sellimonas* abundance was downregulated in rmTBI rats. While, *Rikenellaceae-RC9* and *Ruminococcaceae_UCG-014* abundance showed downregulation in both mild and rmTBI rats.




Figure 5.1a: Alpha-diversity graph.




Figure 5.1b: Beta-diversity graph.





Figure 5.1c: Bacterial abundance.

Figure 5.1a, 5.1b, and 5.1c: showing the change in gut microbiota after 30days of injury.

Metabolic alterations in serum after blunt TBI

We observed significant decrease in formate and lactate at all the time-points after injury. while leucine, isoleucine, valine, alpha-glucose and beta-glucose ratio showed significant decrease at sub- acute and chronic time-point in mild and rmTBI groups. Additionally, the level of NAA was significantly low in both mild and rmTBI groups at all the timepoints except acute mild injury group in compare to control rats.



5.7 Discussion:

The findings of this chapter reveal the metabolic and gut microbiota changes caused by mild and repeated mild diffused closed head injury over time. Diffuse traumatic brain injury (TBI) has a major impact on systemic metabolism, namely cerebral glucose metabolism, and can cause a variety of metabolic and neurological abnormalities. Experimental TBI models in rats have demonstrated unique temporal patterns of brain metabolite alterations at acute, sub acute and chronic timepoints after injury. Along with the previous studies we also observed significant changes in the serum metabolites as early as 4 hr after injury and persisted as long as 30 day after injury. We also found notable change in urine metabolites at 7 day and 30 day PI. Changes in gut microbiome were also observed at 30 day following injury as compared with sham. Significant changes in the GI microbiome were evident after TBI as compared with sham rats, with varying trends among the phylogenetic families.

5.7.1 Systemic Change in serum metabolome following diffused closed head injury:

Serum metabolomics revealed a varied shift in metabolites following mild and repeated mild injuries. Both mild and repetitive mild injuries in rats resulted in decreased energy metabolites, including lactate:pyruvate and $\alpha:\beta$ glucose ratios. Following injury, the branched chain amino acids (BAAs) valine, iso leucine, and leucine were reduced, along with other amino acids such as glutamine, naa, alanine, and taurine, at both the sub-acute and chronic time points. Both injury groups showed a decrease in their $\alpha:\beta$ glucose ratio at sub-acute and chronic time points. Decreased alpha and beta glucose ratio values in serum indicate impaired glucose transport and metabolism, poor glucose transport across the blood-brain barrier, resulting in cerebral metabolic distress and increased mortality following severe brain damage.

Kurtz et al. (2013) conducted a 300-day observational study on 46 consecutive comatose patients with subarachnoid or intracerebral hemorrhage, traumatic brain injury, or cardiac arrest who underwent cerebral microdialysis and intracranial pressure monitoring, and found reduced serum and brain glucose ratios as well as impaired glucose transport. The study also suggests that decreased glucose transport and the blood-brain barrier are related with cerebral metabolic distress and higher mortality following severe brain injury. Another study observed TBI patients for 6 months after injury using intracerebral microdialysis, electroencephalography, and measurements of brain tissue oxygen levels and jugular venous oxygen saturation in 30 patients with traumatic brain injury and reported persistent decrease in extracellular blood glucose. In our study we also observed significant decrease in blood glucose levels following injury at subacute and chronic timepoints. A spike in the lactate-to-pyruvate ratio in serum is a typical metabolic disruption seen in animal investigations. This ratio is a sign of metabolic failure, which is frequently associated with mitochondrial malfunction and hypoxia. A microdialysis investigation on the ballistic-like penetration model of TBI indicated a rise in lactate to pyruvate ratio in rats after 2 hours after injury, and the increase maintained until the 14th day PI.

Another microdialysis study on TBI rats reported significantly higher level of lactate to pyruvate ratio in injured rats after 24hrs of injury. Similarly, we also observed an increase in lactate to pyruvate ratio after injury but significant increase was observed at chronic timepoint only. N-Acetylaspartate (NAA) is a neuron-specific metabolite, and its decrease is a sign of neuronal death. A microdialysis and HPLC investigation on severe TBI patients found an early increase and subsequently decline in NAA levels till 70 hours PI compared to non-TBI individuals. Visser et al. (2023) did a study using magnetic resonance spectroscopy (MRS) or diffusion tensor imaging (DTI) and found a lower level of total NAA in moderate TBI patients at the subacute time point compared to control persons.

Our results also compliment these studies, as we observed the similar change in injured rats at subacute and chronic timepoints following injury. TBI pathophysiology has implications for injury progression and outcome. Identifying metabolic cascades that are altered after acute human mild and

severe TBI, due to breakdown of branched-chain amino acids (BAAs; i.e., valine, isoleucine, and leucine), leads to glucose and energy metabolism, as well as neurotransmitter synthesis and availability. Previous research has also shown that BAA supplementation can help the cognitive recovery of TBI patients (Aquilani et al., 2005).

Jeter et al, 2013 reported decreased levels of all three BAAs in patients with mild TBI relative to healthy volunteers after 24hrs of injury. another study on severe TBI patients reported decreased levels of BAA in plasma after 14days of injury similar to these results we also found significant decrease in BAA levels of injured rat serum.

5.7.2 Gut microbiome changes after mild and rm TBI compare to controls:

The gut microbiome was sequenced in this study, and we discovered that injured rats had lower alpha and beta diversity than sham animals. The structure of a microbial community can be summarized using alpha diversity measures based on its evenness (distribution of group abundances), richness (number of taxonomic groups), or both (Willis, 2019). Alpha metrics that are frequently employed include Shannon's indices (Lemos et al., 2011; Magurran, 2013),

Chao1 (Chao, 1984), observed quantity of amplicon sequence variants (ASVs) (Callahan et al., 2017), and phylogenetic diversity (PD) (Faith, 2006). By taking into account sequence abundances or just the existence or absence of sequences, beta diversity metrics provide a summary of which samples differ from one another. Curtis and Bray, 1957. Alpha diversity was also found to have decreased in TBI rats in previous investigations (Taraskina A et al.; 2022). The phylums of Firmicutes, Bacteroidetes, and Proteobacter showed the maximum alteration. Houlden et al. demonstrated that TBI alters the caecal gut microbiota composition and revealed a connection between the severity of TBI and modifications in the microbiome, particularly in the composition of Firmicutes, Proteobacteria, and Bacteroidetes. The study postulated that these findings might be related to changes in brain function and the stress response brought on by tissue damage. Research revealed that while the phylum Proteobacteria is not

abundant in a healthy human gut, a rise in Proteobacteria indicates gut dysbiosis. In comparison to sham, we also noticed a rise in Proteobacteria following moderate TBI (Shin, Na-Ri et al; 2015). Nicholson et al. (2019) highlighted a significant drop in alpha diversity in moderate TBI rats, as well as a significant decrease in relative abundance at the phylum level in Firmicutes on day 1 after TBI. Specifically, family Lachnospiraceae levels were suppressed after 2 hours to 7 days of PI, although lactobacillace levels showed no significant impact. Our findings also revealed a decrease in Lachnospiraceae levels in TBI rats. Zhanfeng et al. (2022) showed higher Bacteroidetes and lower Firmicutes levels in TBI patients. Bacteroidetes levels were found to be higher in the mild TBI group, while Firmicutes levels were lower in the TBI groups, consistent with previous findings. Furthermore, the mild and rm TBI injured groups showed gut microbiota changes consistent with TBI pathology. A distinguishing feature between mild and rm TBI was increased Proteobacter abundance in mild, which may hinder gut function.

5.8 Conclusion

The present study observed changes in the gut microbiota following injury as well as systemic alterations in metabolism following mild and repeated TBI. Both injury groups revealed a lower alpha:beta glucose ratio and a higher lactate:pyruvate ratio. Signs of oxidative stress and reduced energy metabolites in urine were also indicated by the urine metabolomics of injured rats. Additionally, the gut microbiome showed reduced levels of Firmicutes and Bacteroidetes following injury, as well as altered β diversity and decreased α diversity in rm TBI rats following both traumas. Therefore, the study suggests that while gut dysbiosis and systemic changes are caused by both mild and repeated injury, a different metabolic phenotype with increased inflammation and oxidative stress can be observed by more severe injury.

Chapter 6

Behavioral changes induced by blunt and blast TBI

6.1 Introduction

Traumatic brain injury can induce significant behavioral changes in both humans and animals, particularly at chronic time points after the initial injury. These changes can manifest in a variety of ways, including alterations in mood, cognition, and social functioning (Schretlen & Shapiro, 2003). Previous studies have shown that individuals with traumatic brain injury often experience changes in mood and behavior. For instance, they may exhibit symptoms of depression, such as persistent feelings of sadness or loss of interest in previously enjoyable activities. In addition, individuals with traumatic brain injury may experience heightened levels of anxiety, resulting in increased worry and fear (Li & Liu, 2013). Furthermore, impulsiveness and difficulty with concentration and memory are common behavioral changes observed in both human and animal subjects with traumatic brain injury. These changes in behavior can have a profound impact on the individual's overall quality of life (Dijkers, 2004).

TBI patients are not only at risk for cognitive impairments, but also for developing long-term psychiatric conditions. For example, there is an increased risk for the development of dementia, and even Alzheimer's disease in individuals who have experienced a traumatic brain injury. The exact mechanisms underlying these behavioral changes in traumatic brain injury are not yet fully understood. However, it is believed that the physical damage to the brain, as well as changes in neurotransmitter function and neural connectivity, play a role in these behavioral changes. Furthermore, the social and functional implications of these behavioral changes cannot be ignored. Individuals with traumatic brain injury often experience difficulties in their relationships and social interactions. These behavioral changes can lead to decreased participation in employment and social activities, as well as increased

dependence on others for everyday functional activities. In a study blaze et al. 2020 reported anxiety like behavior in rats after blast induced TBI after 1 to 1.5 months of injury (Blaze et al., 2020). Another TBI study conducted by Arulsamy et al, 2019 using Marmarou's weight drop model reported significant cognitive decline in moderate to severe injury groups after 12 months of injury, although they did not observe any notable change in mild and repetitive mild injury groups at the same time after injury .Although in a lifelong study (24 months) after diffuse blunt TBI on mice reported impaired working memory, decreased spatial memory and vestibule-motor vestibulomotor deficits in rmTBI and mild TBI injury groups compared to sham animals(Arulsamy et al., 2019).

To study the behavioral change in animals we developed two set of injury models based on injury types one is diffused blunt injury and the other is blast induced TBI. Each injury group was divided into three injury group's based on injury frequency i.e control, mild and repetitive mild.

6.2Materials:

Rats

70 % ethanol

Balance beam (2 cm wide × 100 cm long; typically, plastic/polycarbonate with a smooth, nogrip surface that is easily cleaned)

Score sheet

Cotton swabs

6.3Method:

Animal models

Two types of injury groups were created based on injury type (1) Blunt trauma Using modified Marmarou's weight drop model, and (2) Blast trauma using compression driven shock wave tube.

6.3.1Blunt trauma

Three groups—Control, Mild TBI, and rmTBI—were randomly allocated to thirty female Sprague-Dawley (SD) rats weighing 200–250 g. Rats were put on a foam platform and exposed to a free-falling brass rod that was 25 cm above the midpoint of the sagittal plane of the brain for both mild and rmTBI. At 24-hour intervals, the rmTBI group experienced three impacts. Injury induction occurred following a seven-day acclimatization period, and the institutional ethical council authorized all experimental techniques. A regular pellet meal (M/S Golden Feeds, Meharuli, Delhi, India) and unlimited water were given to the rats, who were kept in a 12-hour light/dark cycle at $23\pm1^{\circ}\text{C}$.

6.3.2Blast exposure.

A shock wave was generated using a compression-driven gas shock tube, as previously described (Mishra et al., 2016). Male SD rats were randomly assigned to three groups: sham, mild TBI, and rmTBI, with eight rats in each. Rats were sedated with 3.5% isoflurane and zero air (having approximately 19.9-21.9% oxygen) and put horizontally on a platform within the shockwave tube (Mishra et al., 2016). The mild and moderate TBI groups received a single shock wave dose of 100 kPa. The rmTBI group received three 100 kPa shock wave exposures at 0h, 10h, and 24 hour intervals. The same treatment was performed on sham rats, but they were not exposed to any injuries.

6.4 Behavioral paradigm

All behavioral assessments were administered between 10:00 a.m. and 3:30 p.m. during the light phase of the cycle. Rats were tracked using automated video tracking software (ANYmaze™, Stoelting Co., USA), with the researcher blind to the experimental groups during testing and analysis.

6.4.1 The Neurobehavioral Severity Scale (NSS)

The NSS was used to examine balance, motor coordination (including movement and posture), as well as sensory and motor reflexes at acute (day1), sub-acute (day5), and chronic (day20) stages. Evaluations included balance, landing, tail raise, dragging, righting reflex, ear reflex, eye reflex, sound reflex, tail pinch, and hind paw pinch. The NSS employs a distinct three-point Likert scale, with "0" for normal responses, "1" for incomplete or compromised responses, and "2" for absent responses.

6.4.2 Rotating pole:

The rotating pole test was performed on days 1, 5, and 20 post-injury to assess coordination and movement integration (Ohlsson and Johansson, 1995). The pole was 2.54 cm in diameter and 1.5 m in length, rotating at 4 turns per minute. The pole was divided into three sections, and the time taken for rats to move from one end of the pole to the other was recorded. Falls and reversals were also noted. Rats were trained for two days, followed by a baseline measurement. After injury, three measurements were taken per rat, and the average was calculated.

6.4.3 Open Field Test (OFT)

As previously mentioned, the OFT was used to evaluate emotional, exploratory, and locomotor activity in a new setting (Kumar et al., 2013). The device was 80 cm x 80 cm x 40 cm and was constructed of

plexiglass. A 15-W CFL lamp was positioned 60 cm above the center of the arena floor, which was divided into 16 identical squares. The rats were given five minutes to investigate after being put in a corner that faced the wall. Time spent in the center, average speed, distance traveled, and latency to reach the center were among the behavioral metrics that were noted.

6.4.4 Novel Object Recognition Test (NORT)

In line with the protocol described in an earlier study, the NORT was carried out in the OFT arena (Kumar et al., 2013). There were two stages of the test. In the first, known as the acquisition or learning trial, the rats were given five minutes to investigate two similar objects before going back to their own cage. Following a 5-minute rest, a test trial was conducted where a new object was used in place of a well-known one. The discriminatory ratio was determined by recording the amount of time spent with the known and novel objects during the three-minute test trial.

The Discriminatory Ratio (DR) was calculated as follows:

$$DR = (\text{Total time spent with the novel object}) / (\text{Total time spent with both objects}).$$

6.5 Statistical analysis:

Behavioral changes were compared among the control group, the mild TBI group, and the rmTBI group using a one-way analysis of variance (ANOVA). Graphs and statistical analysis were done using graph pad prism version 5.0 software.

6.6 Results:

6.6.1 Behavioral change after repetitive TBI

NSS

Blunt TBI

Balance, motor coordination, as well as motor and sensory reflexes in rats was observed at acute (day1), sub-acute (day5) and chronic (day20) time-points. We found decreased motor coordination and reflexes in both injury groups in compare to sham. RmTBI rats showed significant change in all the time-points in compare to sham rats. While, mTBI rats showed significant change in acute and sub-acute time-points in compare to sham rats.




Figure 6.1: Showing graph of NSSR experiment following blunt TBI.

Blast TBI

Balance, motor coordination, as well as motor and sensory reflexes in rats was observed at acute, sub-acute and chronic time-points after blast injury. We found decreased motor coordination and reflexes in both injury groups in compare to sham. mTBI rats showed significant change at acute time-point in compare to sham rats. While, rmTBI rats showed significant change in chronic time-point in compare to sham rats.




Figure 6.2: Showing graph of NSSR experiment following blast TBI.

Rotating pole:

Blunt TBI

We observed significant change after rmTBI at acute time point. The coordination and movement were found to decrease in rmTBI rats compare to sham and mildTBI rats




Figure 6.3: Line graph of rotating pole experiment following blunt TBI.

Blast TBI

We observed significant change after rmTBI at all three time-points. The coordination and movement were found to decrease in rmTBI rats compare to sham and mildTBI rats at acute and sub-acute time-point. While, the significant change was observed in rmTBI rats compare to sham rats only.




Figure 6.4: Line graph of rotating pole experiment following blast TBI.

NORT and OFT

Blunt TBI

We observed behavioral change following mild and rmTBI at day30 PI. We found decrease in locomotor activity (total distance travelled and average speed) in mild and rmTBI injury groups respectively compare to sham. While anxiety level was found to increase in mildTBI animals compare to rmTBI injury animals.

NORT results reveled that after injury animals were found to spend more time with novel objects compare to sham, rmTBI groups spent the maximum time with novel object compare to mildTBI and sham group animals. Discriminatory ratio did not show much visible change between injury groups after injury.

We did not observe any significant change in NORT and OFT test of blunt injury groups at chronic time point (day30) PI.




Figure 6.5: Showing the NORT and OFT results following blunt TBI.

Blast TBI

We observed behavioral change following mild and rmTBI at day30 PI. We found significant decrease in locomotor activity (total distance travelled and average speed) in mild and rmTBI injury groups respectively compare to sham. While, anxiety level was found to be in increasing order in mildTBI animals and rmTBI injury animals respectively.

While, blast TBI rats showed significant change in OFT test in mTBI and rmTBI rats at chronic timpoint. Blast TBI rats did not showed any significant change in NORT test.

6.7 Discussion

6.7.1 Alteration in NSS after blast and blunt TBI

NSS is used to evaluate behavioral and sensorimotor effects of TBI on rats that are thought to be the result of changes in neurological functioning (Hamm, 2001; Mahmood et al., 2001; Shohami et al., 1995). We observed significant change in blunt rmTBI rats at acute, sub-acute and chronic timepoints compared to control rats while mTBI rats showed significant changes at acute and sub-acute timepoints compared to control rats. Additionally, the blast TBI rats also showed increase in NSS score at all the timepoints after injury compared to sham but it was significantly upregulated in mTBI rats at acute timepoint and in rmTBI rats at chronic timepoint only. A recent study on moderate to severe closed head injury and decompressive craniectomy model reported increased nss score after injury at day1,

day3, day7, day14 and day28 PI. They also mentioned the significant change in day1, day3, day7 and day28 rats after closed head injury only, compared to sham (Szczygielski et al., 2023). Szczygielski et al, (2016) in a different study reported significant increase in the NSS score in closed head injury rats after 24hrs of injury compared to sham (Szczygielski et al., 2016). Addtitinally, Yarnell et al (2012) reported increased NSS score in mild to moderate blast affected male and female rats at day4 and day8 after injury. Our results also showed the similar changes following TBI (Yarnell et al., 2016).

6.7.2 Alteration in rotating pole after blast and blunt TBI

Roatating pole experiment was performed to evaluate coordination and movement integration ability of rats following TBI (Ohlsson & Johansson, 1995). We found significant decrease in blunt rmTBI only at acute timepoint following injury. While, blast TBI rats showed significant decrease in their score at all three timepoints after TBI. A recent study on closed head injury model reported significant decrease in rotarod score of rmTBI rats after injury (Khan et al., 2023). Mattiason et al (2000) reported significant decrease in the rotating pole score of injured rats at day2 and day7 after FPI injury (Mattiasson et al., 2000). Hoover et al (2004) reported significant decrease in rotating pole scores following FPI rats after day3,1week, 2week, 3week and 4 weeks of injury (Hoover et al., 2004). Divani et al (2015) reported decrease in rotarod scores after injury at 3hrs, 6hrs, 24hrs, 72hrs and 168hrs after blast injury but significant decrease was recorded at 3hrs PI. Therefore, our study is also in line with previous experiments (Divani et al., 2015).

6.7.3 Alteration in cognitive function after blast and blunt TBI

NORT and OFT were done to assess the effect of blunt and blast TBI on the cognitive function of rats. NORT is used to evaluate the memory function of animals following TBI. We did not observe any significant change in NORT after blast and blunt injury at chronic timepoint. OFT or open field test

was done to see the change in locomotor activity and anxiety level of the rats following injury at chronic timepoint. No significant change was found in blunt TBI rats at chronic timepoint. While Blast TBI rats showed significant decrease in their locomotor activity in mild and repetitive mild TBI rats at chronic timepoint. Broussard et al reported significant decrease in OFT (distance travelled) after 5day of rmTBI impact injury mice compared to control (Broussard et al., 2018).

6.8 Conclusion

In conclusion we found that rm and mi TBI significantly affect neurobehavior functions such as NORT, OFT, NSS and rotating pole at acute, sub- acute and chronic timepoints. We observed altered NSS after blunt TBI at all timepoints after injury. While, significant change in rotating pole scores were observed at acute timepoint after blunt rmTBI. After blast injury mi TBI rats showed significant increase in NSS score. While rmTBI rats showed significant change at chronic timepoint. Rotating pole score showed significant decrease in rmTBI rats only after blast induced TBI at all three timepoints. While OFT study showed significant change following blast TBI in both mi and rmTBI groups at chronic timepoint.

Chapter 7

Summary and conclusion

The present thesis entitled Assessment of alterations in neurometabolites and brain microstructure after repetitive mild traumatic brain injury and its association with the behavioral outcome. The present study was intended to explore the histological and metabolic basis of said alterations in the clinical subjects using pre-clinical models. The work under this thesis has focused on TBI which is the leading cause of death and disability in adult population in developing countries. Cerebral and biofluid metabolic profile has been studied to infer a metabolic signature using most reliable/significant set of metabolites altered. Different behavioral aspects of mTBI and rm TBI has been studied.

In chapter 2, NMR metabolomics was used in this study to standardise the classification of diffuse TBI in rats and its effect on the hippocampal metabolome. Ten-week-old SpragueDawley rats were used for the experiments; each group contained 10 animals. A laboratorymade modified Marmarou's weight drop model (Marmarou et al. 1994) was used in this study. TBI was created using a trauma apparatus consisting of a 450-gram metal rod that fell freely through a PVC tube. The brass rod was threaded so that it could be freely lowered from a predetermined height through a 2-metre vertical segment of PVC tubing held in place by an iron support. A stainless-steel button was placed on the anaesthetized animal skull to create a diffuse impact on the brain. The button was placed between the bregma and lambda on the sagittal suture. The rats were placed on a foam bed placed on the base of the iron stand. From a height of 25 cm and 50 cm (mild TBI), 75 cm (moderate TBI), 100 cm, 125 cm, and 150 cm (severe TBI) above the sagittal midline of the rat brain, a 450 gm (1 cm diameter, blunt tip) brass rod was dropped freely. PBS and PFA were employed to anesthetize and perfuse the animals. After a day of wounding and perfusion, the animals were put to death. After being separated, the hippocampus was

liquid nitrogen-snapshot frozen. Tissue samples were then kept for later use at -80 °C. Metabolites were extracted using acetonitrile extraction. Tissues were homogenised in acetonitrile and water (v/v; 1:1). Tissue homogenate was then centrifuged, and the supernatant was collected and lyophilized. For histological studies, whole brains were stored in 4% formalin. Thioflavin S staining was done to observe amyloidosis in the brain after injury. A total of 18 metabolites were identified and observed for metabolic change in the hippocampal tissues after 24 hours of injury. Myo-inositol, choline, taurine, creatine, GABA, glutamate, glutamine, NAA, and lactate were significantly down-regulated after injury. While BAA, acetate, and succinate showed significant upregulation in their levels after injury. An increasing number of beta plaques was observed with injury severity in rats after 24 hours of injury.

In chapter 3, We examined the temporal effects of moderate closed head injury on the hippocampus, thalamus, and striatum. Four groups of five rats each were randomly selected from a total of twenty rats. A modified Marmarou's weight-drop mechanism was used to injure the animals, and samples were taken one, three, and seven days after the injury (PI). After processing the materials, nuclear magnetic resonance (NMR) spectra were acquired for examination. The findings showed metabolic alterations that were specific to time and place. Myo-inositol, creatine, glutamate, succinate, lactate, and N-acetylaspartic acid (NAA) levels in hippocampus tissues changed at 3 days post-infection (PI), whereas tyrosine levels changed at 7 days PI. In the thalamus and striatum, branched-chain amino acids (BAA) and myoinositol levels were altered at 1 day PI. Striatal tissues exhibited significant alterations in taurine, gamma-aminobutyric acid (GABA), choline, and alpha-ketoglutarate (alpha-KG) levels at 1 and 3 days PI. In the thalamus, acetate and GABA levels were altered at 1 day PI, and choline levels were consistently altered at all-time points post-injury. These metabolite alterations may result from disruptions in their associated biochemical pathways. The findings suggest that neurotransmitter and energy metabolism pathways are affected in the hippocampus, thalamus, and striatum following traumatic brain injury (TBI).

In Chapter 4, We investigated metabolic and histological changes in the hippocampus of rats 24 hours after blast-induced damage. Blast-induced traumatic brain injury (BI-TBI) is common among both military personnel and civilians in conflict zones. Individuals in these places may be subjected to repeated blast wave exposures, which can cause cognitive impairments and metabolic abnormalities. The rats were classified into four groups: (i) sham, (ii) mild TBI (mTBI), (iii) moderate TBI (moTBI), and (iv) repetitive mild TBI (rmTBI). Each group experienced varying amounts of blast exposure. Hippocampal tissue samples were taken 24 hours after damage for proton nuclear magnetic resonance spectroscopy (^1H NMR) and immunohistochemistry examination. The metabolic profile of hippocampus tissues indicated significant alterations in glutamate, N-acetylaspartic acid (NAA), acetate, creatine, phosphoethanolamine (PE), ethanolamine, and phosphatidylcholine (PC)/choline levels with these alterations being most pronounced in the rmTBI group. IHC results showed a decrease in acetyl histone 3 (AH3)-positive cells in the rmTBI group compared to other TBI groups and sham rats, indicating potential epigenetic changes due to repeated blast exposure. Furthermore, astrogliosis was noted in the hippocampus of mTBI and moTBI rats, while no significant changes were observed in the rmTBI group.

In chapter 5, The study was aimed to investigate TBI-induced alterations over time, from acute to chronic, following mild and rmTBI. The study used ^1H -NMR metabolomics to examine metabolic changes in tissue, serum, and urine. IHC and faecal 16S rRNA sequencing were used to investigate the potential processes that alter following injury, histological changes, and the gut microbiome. Repetitive mild traumatic brain injury (rmTBI) is a condition in which a person sustains multiple mild traumatic brain injuries over time. Athletes participating in contact sports such as football, hockey, or soccer and persons involved in activities involving a high risk of falls or head impacts are all at risk of repetitive mild TBI. Repeated mild TBIs can have a cumulative effect, which means that each consecutive injury may have a bigger influence on the brain's functioning and healing. A study reported metabolic alterations in collegiate football athletes before and after sports season, and they found alteration in

xanthine metabolism, fatty acid metabolism, medium-chain fatty acid, primary bile acid metabolism and glycolysis, gluconeogenesis, and pyruvate metabolism pathways. The gut microbiota has a close relationship with the central nervous system (CNS). Studies have demonstrated that traumatic brain injury can alter the diversity and abundance of gut microbial species, leading to dysbiosis and the development of virulent phylogenetic changes within the intestinal commensal microbiota. This disruption of the gut microbiome following a traumatic brain injury is thought to be a result of the bidirectional communication between the brain and the gut. Thirty female SD rats weighing 200 to 250 g were randomly assigned to one of three groups: control, mild TBI, or rmTBI. For mild and rmTBI, rats were placed on a foam bed and subjected to a free-falling brass rod was dropped from a height of 25 cm, on the sagittal midway of the animal skull. Rm TBI rats were subjected to three impacts at 24-hour intervals. Serum

(4 hours, 5 days, and 30 days) and faecal samples (30 days) were collected for NMR and 16Sr RNA analysis. The α -diversity showed an increase in richness within the mild group, while the rmTBI group showed a decrease in richness. At the phylum level, the firmicutes and bacteria decreased in the rmTBI group. While phylum firmicutes was downregulated in mild TBI rats compared to sham, Other bacterial phylum, including Spirochaetes, Proteobacteria, and Patescibacteria, were also altered after injury. Mildly injured rats showed increased proteobacteria abundance. While serum metabolites ratio of alpha and beta glucose showed a significant increase at the sub-acute and chronic time- points in the mild and rmTBI groups. BAA showed decreases at sub-acute and chronic time- points in the mild and rmTBI groups. Lactate and pyruvate ratio levels were upregulated at chronic timepoints in both mild and repeated mild rats. Additionally, there was a significant decrease in VLDL/LDL at all the time points except for the for the mild 4 hours after injury.

In chapter 6, we focused in the TBI induced significant behavioural changes in rodents, particularly at chronic time points after the initial injury. These changes can manifest in a variety of ways, including alterations in mood, cognition, and social functioning. Research has shown that individuals with

traumatic brain injuries often experience changes in mood and behavior. To study the behavioural change in animals, we developed two sets of injury models based on injury types: one is diffused blunt injury, and the other is blast-induced BI. Each injury group was divided into three groups based on severity, i.e., control, mild, and repetitive mild. Thirty female SD rats weighing 200 to 250 g were randomly assigned to one of three groups: control, mild TBI, or rmTBI. For mild and rmTBI, rats were placed on a foam bed and subjected to a free-falling brass rod from a height of 25 cm, above the sagittal halfway point of the rat brain. Rm TBI rats were subjected to three impacts at 24-hour intervals. In the case of blunt TBI NSSR scoring, we observed decreased motor coordination and reflexes in both injury groups in comparison to sham. RmTBI rats showed a significant change in all time points compared to sham rats. While mTBI rats showed significant changes in acute and sub-acute time points in comparison to sham rats, while in the rotating pole experiment, blunt TBI rats showed significant change after rmTBI at the acute time point. The coordination and movement were found to decrease in rmTBI rats compared to sham and mildTBI rats. Additionally, we did not observe any significant changes in the NORT and OFT tests after blunt trauma.

On the other hand, in blast TBI rats, we found decreased motor coordination and reflexes in both injury groups compared to sham. mTBI rats showed a significant change at the acute time point in comparison to sham rats. While rmTBI rats showed a significant change in chronic time points compared to sham rats, In the rotating pole experiment, we observed a significant change after rmTBI at all three time points. The coordination and movement were found to decrease in rmTBI rats compared to sham and mildTBI rats at acute and sub-acute time- points. While a significant change was observed in rmTBI rats compared to sham rats only, We observed behavioural change following mild and rmTBI at day 30 PI. We found a significant decrease in locomotor activity (total distance travelled and average speed) (OFT) in the mild and rmTBI injury groups, respectively, compared to sham. While anxiety levels were found to be in increasing order in mildTBI animals and rmTBI injury animals, respectively, The NORT study (time spent with novel objects and discriminatory ratio) results showed a decrease in the time

spent with novel objects and discriminatory ratio in injury animals compared to sham at day 30 PI. Although this decrease was not significant.

The highlights of the thesis are:

1. Increasing height leads to increased mortality in rats with impaired energy metabolites and neurite plaques.
2. Blunt impact to the brain leads to regional variation in energy metabolites resulting in dysregulated TCA cycle.
3. Repetitive Blast exposure is accompanied with alterations in the acetate and NAA levels leading to decreased acetylation.
4. At systemic level, Blunt impact leads to equivalent damage in both mild and repeated mild injury. While blast exposure has variable effect after mild and repeated mild exposure.
5. Glycolysis and TCA pathways intermediates are majorly affected in TBI and seen in serum metabolomic analysis.

References

Akram, M. (2014). Citric acid cycle and role of its intermediates in metabolism. *Cell Biochemistry and Biophysics*, 68(3), 475–478. <https://doi.org/10.1007/s12013-013-9750-1>

Albrecht, J., Sidoryk-Węgrzynowicz, M., Zielińska, M., & Aschner, M. (2010). Roles of glutamine in neurotransmission. *Neuron Glia Biology*, 6(4), 263–276. <https://doi.org/10.1017/S1740925X11000093>

Álvarez, Z., Castaño, O., Castells, A. A., Mateos-Timoneda, M. A., Planell, J. A., Engel, E., & Alcántara, S. (2014). Neurogenesis and vascularization of the damaged brain using a lactatereleasing biomimetic scaffold. *Biomaterials*, 35(17), 4769–4781. <https://doi.org/10.1016/j.biomaterials.2014.02.051>

Andersen, J. V., Skotte, N. H., Aldana, B. I., Nørremølle, A., & Waagepetersen, H. S. (2019). Enhanced cerebral branched-chain amino acid metabolism in R6/2 mouse model of Huntington's disease. *Cellular and Molecular Life Sciences*, 76(12), 2449–2461. <https://doi.org/10.1007/s00018-019-03051-2>

Aquilani, R., Iadarola, P., Contardi, A., Boselli, M., Verri, M., Pastorini, O., Boschi, F., Arcidiaco, P., & Viglio, S. (2005). Branched-chain amino acids enhance the cognitive recovery of patients with severe traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*, 86(9), 1729–1735. <https://doi.org/10.1016/j.apmr.2005.03.022>

Arcelli, P., Frassoni, C., Regondi, M. C., De Biasi, S., & Spreafico, R. (1997). GABAergic neurons in mammalian thalamus: A marker of thalamic complexity? *Brain Research Bulletin*, 42(1), 27–37. [https://doi.org/10.1016/s0361-9230\(96\)00107-4](https://doi.org/10.1016/s0361-9230(96)00107-4)

Arulsamy, A., Corrigan, F., & Collins-Praino, L. E. (2019). Cognitive and neuropsychiatric impairments vary as a function of injury severity at 12 months post-experimental diffuse traumatic brain injury: Implications for dementia development. *Behavioural Brain Research*, 365, 66–76. <https://doi.org/10.1016/j.bbr.2019.02.045>

Andres, R. H., Ducray, A. D., Schlattner, U., Wallimann, T., & Widmer, H. R. (2008). Functions and effects of creatine in the central nervous system. *Brain Research Bulletin*, 76(4), 329–343. <https://doi.org/10.1016/j.brainresbull.2008.02.035>

Arciniegas, D. B., Held, K., & Wagner, P. (2002). Cognitive Impairment following TBI. Current treatment options in neurology, 4(1), 43–57.

Ariyannur, P. S., Moffett, J. R., Madhavarao, C. N., Arun, P., Vishnu, N., Jacobowitz, D. M., et al. (2010). Nuclear-cytoplasmic localization of acetyl coenzyme a synthetase-1 in the rat brain, 518(15), 2952–2977. <https://doi.org/10.1002/cne.22373>. Nuclear-cytoplasmic

Attwell, D., Buchan, A. M., Charpak, S., Lauritzen, M., MacVicar, B. A., & Newman, E. A. (2010). Glial and neuronal control of brain blood flow. *Nature*, 468(7321), 232–243. <https://doi.org/10.1038/nature09613>

Barkai, A. I., Dunner, D. L., Gross, H. A., Mayo, P., & Fieve, R. R. (1978). Reduced myo-inositol levels in cerebrospinal fluid from patients with affective disorder. *Biological Psychiatry*, 13(1), 65–72.

Blaze, J., Choi, I., Wang, Z., Umali, M., Mendelev, N., Tschiffely, A. E., Ahlers, S. T., Elder, G. A., Ge, Y., & Haghghi, F. (2020). Blast-Related Mild TBI Alters Anxiety-Like Behavior and Transcriptional Signatures in the Rat Amygdala. *Frontiers in Behavioral Neuroscience*, 14, 160. <https://doi.org/10.3389/fnbeh.2020.00160>

Briend, F., Nelson, E. A., Maximo, O., Armstrong, W. P., Kraguljac, N. V., & Lahti, A. C. (2020). Hippocampal glutamate and hippocampus subfield volumes in antipsychotic-naïve first episode psychosis subjects and relationships to duration of untreated psychosis. *Translational Psychiatry*, 10(1), 137. <https://doi.org/10.1038/s41398-020-0812-z>

Banoei, M. M., Casault, C., Metwaly, S. M., & Winston, B. W. (2018). Metabolomics and Biomarker Discovery in Traumatic Brain Injury. *Journal of Neurotrauma*, 35(16), 1831–1848. <https://doi.org/10.1089/neu.2017.5326>

Brosnan, J. T., & Brosnan, M. E. (2007). Creatine: Endogenous Metabolite, Dietary, and Therapeutic Supplement. Annual Review of Nutrition, 27(1), 241–261. <https://doi.org/10.1146/annurev.nutr.27.061406.093621>

Broussard, J. I., Acion, L., De Jesús-Cortés, H., Yin, T., Britt, J. K., Salas, R., Costa-Mattioli, M., Robertson, C., Pieper, A. A., Arciniegas, D. B., & Jorge, R. (2018). Repeated mild traumatic brain injury produces neuroinflammation, anxiety-like behaviour and impaired spatial memory in mice. *Brain Injury*, 32(1), 113–122. <https://doi.org/10.1080/02699052.2017.1380228>

Calvetti, D., & Somersalo, E. (2013). Quantitative in silico Analysis of Neurotransmitter Pathways Under Steady State Conditions. *Frontiers in Endocrinology*, 4(OCT), 1–14.
<https://doi.org/10.3389/fendo.2013.00137>

Caplan, B., Bogner, J., Brenner, L., Malec, J., Sharma, B., Lawrence, D. W., & Hutchison, M. G. (2018). Branched chain amino acids (BCAAs) and traumatic brain injury: a systematic review. *Journal of head trauma rehabilitation*, 33(1), 33–45.

Carpenter, K. L. H., Jalloh, I., & Hutchinson, P. J. (2015). Glycolysis and the significance of lactate in traumatic brain injury. *Frontiers in Neuroscience*, 9(APR), 1–15.
<https://doi.org/10.3389/fnins.2015.00112>

Clark, A. L., Merritt, V. C., Bigler, E. D., Bangen, K. J., Werhane, M., Sorg, S. F., et al. (2018). Blast-exposed veterans with mild traumatic brain injury show greater frontal cortical thinning and poorer executive functioning. *Frontiers in Neurology*, 9(NOV), 1–11.
<https://doi.org/10.3389/fneur.2018.00873>

Clark, J. F., Doepke, A., Filosa, J. A., Wardle, R. L., Lu, A., Meeker, T. J., & Pyne-Geithman, G. J. (2006). N-Acetylaspartate as a reservoir for glutamate. *Medical Hypotheses*, 67(3), 506–512. <https://doi.org/10.1016/j.mehy.2006.02.047>

Carron, S. F., Sun, M., Shultz, S. R., & Rajan, R. (2020). Inhibitory neuronal changes following a mixed diffuse-focal model of traumatic brain injury. *Journal of Comparative Neurology*, 528(2), 175–198. <https://doi.org/10.1002/cne.24746>

Clark, A. L., Merritt, V. C., Bigler, E. D., Bangen, K. J., Werhane, M., Sorg, S. F., Bondi, M. W., Schiehser, D. M., & Delano-Wood, L. (2018). Blast-exposed veterans with mild traumatic brain injury show greater frontal cortical thinning and poorer executive functioning. *Frontiers in Neurology*, 9(NOV), 1–11.
<https://doi.org/10.3389/fneur.2018.00873>

Daubner, S. C., Le, T., & Wang, S. (2011). Tyrosine hydroxylase and regulation of dopamine synthesis. *Archives of Biochemistry and Biophysics*, 508(1), 1–12.
<https://doi.org/10.1016/j.abb.2010.12.017>

Deutch, A. Y., Colbran, R. J., & Winder, D. J. (2007). Striatal plasticity and medium spiny neuron dendritic remodeling in parkinsonism. *Parkinsonism & Related Disorders*, 13, S251–S258.
[https://doi.org/10.1016/S1353-8020\(08\)70012-9](https://doi.org/10.1016/S1353-8020(08)70012-9)

Di Pietro, V., Amorini, A. M., Tavazzi, B., Vagnozzi, R., Logan, A., Lazzarino, G., Signoretti, S., Lazzarino, G., & Belli, A. (2014). The Molecular Mechanisms Affecting N-Acetylaspartate Homeostasis Following Experimental Graded Traumatic Brain Injury. *Molecular Medicine*, 20(1), 147–157. <https://doi.org/10.2119/molmed.2013.00153>

D'Adamo Jr, A. F., Smith, J. C., & Woiler, C. (1972). The occurrence of N-acetylaspartate amidohydrolase (Aminoacylase n) in the developing rat. *Journal of Neurochemistry*, 20(4), 1275–1278. [https://doi.org/10.1016/0032-5910\(72\)80020-2](https://doi.org/10.1016/0032-5910(72)80020-2)

Dedeoglu, A., Choi, J.-K., Cormier, K., Kowall, N. W., & Jenkins, B. G. (2004). Magnetic resonance spectroscopic analysis of Alzheimer's disease mouse brain that express mutant human APP shows altered neurochemical profile. *Brain Research*, 1012(1–2), 60–65. <https://doi.org/10.1016/j.brainres.2004.02.079>

Dienel, G. A. (2014). Lactate shuttling and lactate use as fuel after traumatic brain injury: metabolic considerations. *Journal of Cerebral Blood Flow & Metabolism*, 34(11), 1736–1748.

Dijkers, M. P. (2004). Quality of life after traumatic brain injury: A review of research approaches and findings¹¹No commercial party having a direct financial interest in the results of the research supporting this article has or will confer a benefit upon the authors(s) or upon any organization with which the author(s) is/are associated. *Archives of Physical Medicine and Rehabilitation*, 85, 21–35. <https://doi.org/10.1016/j.apmr.2003.08.119>

Divani, A. A., Murphy, A. J., Meints, J., Sadeghi-Bazargani, H., Nordberg, J., Monga, M., Low, W. C., Bhatia, P. M., Beilman, G. J., & SantaCruz, K. S. (2015). A Novel Preclinical Model of Moderate Primary Blast-Induced Traumatic Brain Injury. *Journal of Neurotrauma*, 32(14), 1109–1116. <https://doi.org/10.1089/neu.2014.3686>

Estrada Sánchez, A. M., Mejía-Toiber, J., & Massieu, L. (2008). Excitotoxic Neuronal Death and the Pathogenesis of Huntington's Disease. *Archives of Medical Research*, 39(3), 265–276. <https://doi.org/10.1016/j.arcmed.2007.11.011>

Fernstrom, J. D., & Fernstrom, M. H. (2007). Tyrosine, Phenylalanine, and Catecholamine Synthesis and Function in the Brain ,2,. *The Journal of Nutrition*, 137(6), 1539S–1547S. <https://doi.org/10.1093/jn/137.6.1539S>

Frankowski, J. C., Tierno, A., Pavani, S., Cao, Q., Lyon, D. C., & Hunt, R. F. (2022). Brain-wide reconstruction of inhibitory circuits after traumatic brain injury. *Nature Communications*, 13(1), 3417. <https://doi.org/10.1038/s41467-022-31072-2>

Fromm, L., Heath, D. L., Vink, R., & Nimmo, A. J. (2004). Magnesium attenuates post-traumatic depression/anxiety following diffuse traumatic brain injury in rats. *Journal of the American College of Nutrition*, 23(5), 529S-533S. <https://doi.org/10.1080/07315724.2004.10719396>

Gama Sosa, M. A., De Gasperi, R., Perez Garcia, G. S., Perez, G. M., Searcy, C., Vargas, D., Spencer, A., Janssen, P. L., Tschiffely, A. E., McCarron, R. M., Ache, B., Manoharan, R., Janssen, W. G., Tappan, S. J., Hanson, R. W., Gandy, S., Hof, P. R., Ahlers, S. T., & Elder, G. A. (2019). Low-level blast exposure disrupts gliovascular and neurovascular connections and induces a chronic vascular pathology in rat brain. *Acta Neuropathologica Communications*, 7(1), 6. <https://doi.org/10.1186/s40478-018-0647-5>

Gama Sosa, M. A., De Gasperi, R., Perez Garcia, G. S., Perez, G. M., Searcy, C., Vargas, D., et al. (2019). Low-level blast exposure disrupts gliovascular and neurovascular connections and induces a chronic vascular pathology in rat brain. *Acta Neuropathologica Communications*, 7(1), 6. <https://doi.org/10.1186/s40478-018-0647-5>

Gao, W.-M., Chadha, M. S., Kline, A. E., Clark, R. S. B., Kochanek, P. M., Dixon, C. E., & Jenkins, L. W. (2006). Immunohistochemical analysis of histone H3 acetylation and methylation—Evidence for altered epigenetic signaling following traumatic brain injury in immature rats. *Brain Research*, 1070(1), 31–34. <https://doi.org/10.1016/j.brainres.2005.11.038>

Ge, S., Pradhan, D. A., Ming, G., & Song, H. (2007). GABA sets the tempo for activity-dependent adult neurogenesis. *Trends in Neurosciences*, 30(1), 1–8. <https://doi.org/10.1016/j.tins.2006.11.001>

Guilfoyle, M. R., Helmy, A., Donnelly, J., Stovell, M. G., Timofeev, I., Pickard, J. D., Czosnyka, M., Smielewski, P., Menon, D. K., Carpenter, K. L. H., & Hutchinson, P. J. (2021). Characterising the dynamics of cerebral metabolic dysfunction following traumatic brain injury: A microdialysis study in 619 patients. *PLOS ONE*, 16(12), e0260291. <https://doi.org/10.1371/journal.pone.0260291>

Hamm, R. J. (2001). Neurobehavioral Assessment of Outcome following Traumatic Brain Injury in Rats: An Evaluation of Selected Measures. *Journal of Neurotrauma*, 18(11), 1207–1216.
<https://doi.org/10.1089/089771501317095241>

Harvey, B. H., Brand, L., Jeeva, Z., & Stein, D. J. (2006). Cortical/hippocampal monoamines, HPAaxis changes and aversive behavior following stress and restress in an animal model of posttraumatic stress disorder. *Physiology & Behavior*, 87(5), 881–890.
<https://doi.org/10.1016/j.physbeh.2006.01.033>

Hoge, C. W., McGurk, D., Thomas, J. L., Cox, A. L., Engel, C. C., & Castro, C. A. (2015). Mild Traumatic Brain Injury in U.S. Soldiers Returning from Iraq Charles. *New England Journal of Medicine*, 358(5), 453–463.

Hoover, R. C., Motta, M., Davis, J., Saatman, K. E., Fujimoto, S. T., Thompson, H. J., Stover, J. F., Dichter, M. A., Twyman, R., White, H. S., & McIntosh, T. K. (2004). Differential Effects of the Anticonvulsant Topiramate on Neurobehavioral and Histological Outcomes following Traumatic Brain Injury in Rats. *Journal of Neurotrauma*, 21(5), 501–512.
<https://doi.org/10.1089/089771504774129847>

Huerta-Ocampo, I., Hacioglu-Bay, H., Dautan, D., & Mena-Segovia, J. (2020). Distribution of Midbrain Cholinergic Axons in the Thalamus. *Eneuro*, 7(1), ENEURO.0454-19.2019.
<https://doi.org/10.1523/ENEURO.0454-19.2019>

Ikarashi, Y., Takahashi, A., Ishimaru, H., Arai, T., & Maruyama, Y. (2002). Relations Between the Extracellular Concentrations of Choline and Acetylcholine in Rat Striatum. *Journal of Neurochemistry*, 69(3), 1246–1251. <https://doi.org/10.1046/j.1471-4159.1997.69031246.x>

Jaeger, C., Glaab, E., Michelucci, A., Binz, T. M., Koeglsberger, S., Garcia, P., Trezzi, J.-P., Ghelfi, J., Balling, R., & Buttini, M. (2015). The Mouse Brain Metabolome. *The American Journal of Pathology*, 185(6), 1699–1712. <https://doi.org/10.1016/j.ajpath.2015.02.016>

Jager, P., Moore, G., Calpin, P., Durmishi, X., Kita, Y., Salgarella, I., Wang, Y., Schultz, S. R., Brickley, S., Shimogori, T., & Delogu, A. (2019). *Dual midbrain and forebrain origins of thalamic inhibitory interneurons* [Preprint]. *Developmental Biology*.
<https://doi.org/10.1101/651745>

Jeter, C. B., Hergenroeder, G. W., Ward, N. H., Moore, A. N., & Dash, P. K. (2013). Human Mild Traumatic Brain Injury Decreases Circulating Branched-Chain Amino Acids and Their Metabolite Levels. *Journal of Neurotrauma*, 30(8), 671–679.
<https://doi.org/10.1089/neu.2012.2491>

Kamnaksh, A., Ahmed, F., Kovesdi, E., Barry, E. S., Grunberg, N. E., Long, J. B., & Agoston, D. V. (2014). Molecular mechanisms of increased cerebral vulnerability after repeated mild blastinduced traumatic brain injury. *Translational Proteomics*, 3, 22–37.
<https://doi.org/10.1016/j.trprot.2013.11.001>

Kelsom, C., & Lu, W. (2013). Development and specification of GABAergic cortical interneurons. *Cell & Bioscience*, 3(1), 19. <https://doi.org/10.1186/2045-3701-3-19>

Khan, A. R., Zehra, S., Baranwal, A. K., Kumar, D., Ali, R., Javed, S., & Bhaisora, K. (2023). Whole-Blood Metabolomics of a Rat Model of Repetitive Concussion. *Journal of Molecular Neuroscience*, 73(9–10), 843–852. <https://doi.org/10.1007/s12031-023-02162-7>

Kontro, P. (1987). Interactions of Taurine and Dopamine in the Striatum. In R. J. Huxtable, F. Franconi, & A. Giotti (Eds.), *The Biology of Taurine* (Vol. 217, pp. 347–355). Springer US.
https://doi.org/10.1007/978-1-4899-0405-8_37

Kumari, M., Arora, P., Sharma, P., Hasija, Y., Rana, P., D'souza, M. M., Chandra, N., & Trivedi, R. (2023). Acute metabolic alterations in the hippocampus are associated with decreased acetylation after blast induced TBI. *Metabolomics*, 19(1), 5. <https://doi.org/10.1007/s11306-022-01970-z>

Kwok, Y. N., & Collier, B. (1982). Synthesis of Acetylcholine from Acetate in a Sympathetic Ganglion. *Journal of Neurochemistry*, 39(1), 16–26. <https://doi.org/10.1111/j.1471-4159.1982.tb04696.x>

Leem, Y.-H., Kato, M., & Chang, H. (2018). Regular exercise and creatine supplementation prevent chronic mild stress-induced decrease in hippocampal neurogenesis via Wnt/GSK3 β /β-catenin pathway. *Journal of Exercise Nutrition & Biochemistry*, 22(2), 1–6.
<https://doi.org/10.20463/jenb.2018.0009>

Li, L., & Liu, J. (2013). The effect of pediatric traumatic brain injury on behavioral outcomes: A systematic review. *Developmental Medicine & Child Neurology*, 55(1), 37–45.
<https://doi.org/10.1111/j.1469-8749.2012.04414.x>

Li, Teng, En Hu, Pengfei Li, Zhaoyu Yang, Yao Wu, Ruoqi Ding, Xiaofei Zhu, Tao Tang, and Yang Wang. "Metabolomics deciphers potential targets of Xuefu Zhuyu decoction against traumatic brain injury in rat." *Frontiers in Pharmacology* 11 (2020): 559618.

Lim, S. A. O., Kang, U. J., & McGehee, D. S. (2014). Striatal cholinergic interneuron regulation and circuit effects. *Frontiers in Synaptic Neuroscience*, 6, 22.
<https://doi.org/10.3389/fnsyn.2014.00022>

Liu, S., He, L., & Yao, K. (2018). The Antioxidative Function of Alpha-Ketoglutarate and Its Applications. *BioMed Research International*, 2018, 3408467.
<https://doi.org/10.1155/2018/3408467>

Maffezzini, C., Calvo-Garrido, J., Wredenberg, A., & Freyer, C. (2020). Metabolic regulation of neurodifferentiation in the adult brain. *Cellular and Molecular Life Sciences: CMLS*, 77(13), 2483–2496. <https://doi.org/10.1007/s00018-019-03430-9>

Mahmood, A., Lu, D., Wang, L., Li, Y., Lu, M., & Chopp, M. (2001). Treatment of traumatic brain injury in female rats with intravenous administration of bone marrow stromal cells. *Neurosurgery*, 49(5), 1196–1203; discussion 1203-1204.

Martínez-Reyes, I., & Chandel, N. S. (2020). Mitochondrial TCA cycle metabolites control physiology and disease. *Nature Communications*, 11(1), 102. <https://doi.org/10.1038/s41467019-13668-3>

Matsui, T., Omuro, H., Liu, Y.-F., Soya, M., Shima, T., McEwen, B. S., & Soya, H. (2017). Astrocytic glycogen-derived lactate fuels the brain during exhaustive exercise to maintain endurance capacity. *Proceedings of the National Academy of Sciences*, 114(24), 6358–6363.
<https://doi.org/10.1073/pnas.1702739114>

Mattiasson, G. J., Philips, M. F., Tomasevic, G., Johansson, B. B., Wieloch, T., & McIntosh, T. K. (2000). The rotating pole test: Evaluation of its effectiveness in assessing functional motor deficits following

experimental head injury in the rat. *Journal of Neuroscience Methods*, 95(1), 75–82.
[https://doi.org/10.1016/S0165-0270\(99\)00162-4](https://doi.org/10.1016/S0165-0270(99)00162-4)

McGeer, P. L., McGeer, E. G., & Nagai, T. (1983). GABAergic and cholinergic indices in various regions of rat brain after intracerebral injections of folic acid. *Brain Research*, 260(1), 107–116. [https://doi.org/10.1016/0006-8993\(83\)90768-0](https://doi.org/10.1016/0006-8993(83)90768-0)

McGuire, J. L., DePasquale, E. A. K., Watanabe, M., Anwar, F., Ngwenya, L. B., Atluri, G., RomickRosendale, L. E., McCullumsmith, R. E., & Evanson, N. K. (2019). Chronic Dysregulation of Cortical and Subcortical Metabolism After Experimental Traumatic Brain Injury. *Molecular Neurobiology*, 56(4), 2908–2921. <https://doi.org/10.1007/s12035-018-1276-5>

McGuire, J. L., Ngwenya, L. B., & McCullumsmith, R. E. (2018). Neurotransmitter changes after traumatic brain injury: An update for new treatment strategies. *Molecular Psychiatry*, 24(7), 995–1012. <https://doi.org/10.1038/s41380-018-0239-6>

McKenna, M. C. (2007). The glutamate-glutamine cycle is not stoichiometric: Fates of glutamate in brain. *Journal of Neuroscience Research*, 85(15), 3347–3358.
<https://doi.org/10.1002/jnr.21444>

Mishkovsky, M., Comment, A., & Gruetter, R. (2012). *In Vivo* Detection of Brain Krebs Cycle Intermediate by Hyperpolarized Magnetic Resonance. *Journal of Cerebral Blood Flow & Metabolism*, 32(12), 2108–2113. <https://doi.org/10.1038/jcbfm.2012.136>

Moffett, J. R., Arun, P., Ariyannur, P. S., & Namboodiri, A. M. A. (2013). N-Acetylaspartate reductions in brain injury: Impact on post-injury neuroenergetics, lipid synthesis, and protein acetylation. *Frontiers in Neuroenergetics*, 5, 11. <https://doi.org/10.3389/fnene.2013.00011>

Morland, C., Andersson, K. A., Haugen, Ø. P., Hadzic, A., Kleppa, L., Gille, A., Rinholm, J. E., Palibrk, V., Diget, E. H., Kennedy, L. H., Stølen, T., Hennestad, E., Moldestad, O., Cai, Y., Puchades, M., Offermanns, S., Vervaeke, K., Bjørås, M., Wisløff, U., ... Bergersen, L. H. (2017). Exercise induces cerebral VEGF and angiogenesis via the lactate receptor HCAR1. *Nature Communications*, 8(1), 15557. <https://doi.org/10.1038/ncomms15557>

Moujahid, A., D'Anjou, A., & GraÃ±a, M. (2014). Energy demands of diverse spiking cells from the neocortex, hippocampus, and thalamus. *Frontiers in Computational Neuroscience*, 8.

<https://doi.org/10.3389/fncom.2014.00041>

Nakatomi, H., Kuriu, T., Okabe, S., Yamamoto, S., Hatano, O., Kawahara, N., Tamura, A., Kirino, T., & Nakafuku, M. (2002). Regeneration of Hippocampal Pyramidal Neurons after Ischemic Brain Injury by Recruitment of Endogenous Neural Progenitors. *Cell*, 110(4), 429–441.

[https://doi.org/10.1016/S0092-8674\(02\)00862-0](https://doi.org/10.1016/S0092-8674(02)00862-0)

Oh, S. J., Lee, H.-J., Jeong, Y. J., Nam, K. R., Kang, K. J., Han, S. J., Lee, K. C., Lee, Y. J., & Choi, J. Y. (2020). Evaluation of the neuroprotective effect of taurine in Alzheimer's disease using functional molecular imaging. *Scientific Reports*, 10(1), 15551.

<https://doi.org/10.1038/s41598-020-72755-4>

Ohlsson, A.-L., & Johansson, B. B. (1995). Environment Influences Functional Outcome of Cerebral Infarction in Rats. *Stroke*, 26(4), 644–649. <https://doi.org/10.1161/01.STR.26.4.644>

Peskind, E. R., Petrie, E. C., Cross, D. J., Pagulayan, K., McCraw, K., Hoff, D., Hart, K., Yu, C.-E., Raskind, M. A., Cook, D. G., & Minoshima, S. (2011). Cerebrocerebellar hypometabolism associated with repetitive blast exposure mild traumatic brain injury in 12 Iraq war Veterans with persistent post-concussive symptoms. *NeuroImage*, 54(SUPPL. 1), S76–S82.

<https://doi.org/10.1016/j.neuroimage.2010.04.008>

Piao, C.-S., Holloway, A. L., Hong-Routson, S., & Wainwright, M. S. (2019). Depression following traumatic brain injury in mice is associated with down-regulation of hippocampal astrocyte glutamate transporters by thrombin. *Journal of Cerebral Blood Flow and Metabolism: Official Journal of the International Society of Cerebral Blood Flow and Metabolism*, 39(1), 58–73. <https://doi.org/10.1177/0271678X17742792>

Prins, M., Alexander, D., Giza, C. C., & Hovda, D. A. (2013). Repeated Mild Traumatic Brain Injury: Mechanisms of Cerebral Vulnerability. *The Journal of Neurotrauma*, 30(1), 30–38.

<https://doi.org/10.1089/neu.2012.2399>

Rana, P., Rama Rao, K. V., Ravula, A., Trivedi, R., D'Souza, M., Singh, A. K., Gupta, R. K., & Chandra, N. (2020a). Oxidative stress contributes to cerebral metabolomic profile changes in animal model of blast-induced traumatic brain injury. *Metabolomics*, 16(3), 39.

<https://doi.org/10.1007/s11306-020-1649-4>

Rana, P., Rama Rao, K. V., Ravula, A., Trivedi, R., D'Souza, M., Singh, A. K., Gupta, R. K., & Chandra, N. (2020b). Oxidative stress contributes to cerebral metabolomic profile changes in animal model of blast-induced traumatic brain injury. *Metabolomics*, 16(3), 1–12.

<https://doi.org/10.1007/s11306-020-1649-4>

Rivas-Arancibia, S., Rodríguez, A. I., Zigova, T., Willing, A. E., Brown, W. D., Cahill, D. W., & Sanberg, P. R. (2001). Taurine increases rat survival and reduces striatal damage caused by 3nitropropionic acid. *The International Journal of Neuroscience*, 108(1–2), 55–67.

<https://doi.org/10.3109/00207450108986505>

Ruotsalainen, M., & Ahtee, L. (1996). Intrastriatal taurine increases striatal extracellular dopamine in a tetrodotoxin-sensitive manner in rats. *Neuroscience Letters*, 212(3), 175–178.

[https://doi.org/10.1016/0304-3940\(96\)12821-4](https://doi.org/10.1016/0304-3940(96)12821-4)

Saija, A., Hayes, R. L., Lyeth, B. G., Edward Dixon, C., Yamamoto, T., & Robinson, S. E. (1988).

The effect of concussive head injury on central cholinergic neurons. *Brain Research*, 452(1–2), 303–311. [https://doi.org/10.1016/0006-8993\(88\)90034-0](https://doi.org/10.1016/0006-8993(88)90034-0)

Schlett, K. (2006). Glutamate as a Modulator of Embryonic and Adult Neurogenesis. *Current Topics in Medicinal Chemistry*, 6(10), 949–960. <https://doi.org/10.2174/156802606777323665>

Schretlen, D. J., & Shapiro, A. M. (2003). A quantitative review of the effects of traumatic brain injury on cognitive functioning. *International Review of Psychiatry*, 15(4), 341–349.

<https://doi.org/10.1080/09540260310001606728>

Scremin, O. U., Li, M. G., Roch, M., Booth, R., & Jenden, D. J. (2006). Acetylcholine and choline dynamics provide early and late markers of traumatic brain injury. *Brain Research*, 1124(1), 155–166. <https://doi.org/10.1016/j.brainres.2006.09.062>

Shank, R. P., & Campbell, G. L. (1982). Glutamine and alpha-ketoglutarate uptake and metabolism by nerve terminal enriched material from mouse cerebellum. *Neurochemical Research*, 7(5),

601–616. <https://doi.org/10.1007/BF00965126>

Shin, S. S., & Dixon, C. E. (2015). Alterations in Cholinergic Pathways and Therapeutic Strategies Targeting Cholinergic System after Traumatic Brain Injury. *Journal of Neurotrauma*, 32(19), 1429–1440. <https://doi.org/10.1089/neu.2014.3445>

Shirayama, Y., Takahashi, M., Osone, F., Hara, A., & Okubo, T. (2017). Myo-inositol, Glutamate, and Glutamine in the Prefrontal Cortex, Hippocampus, and Amygdala in Major Depression. *Biological Psychiatry. Cognitive Neuroscience and Neuroimaging*, 2(2), 196–204.

<https://doi.org/10.1016/j.bpsc.2016.11.006>

Shohami, E., Novikov, M., & Bass, R. (1995). Long-term effect of HU-211, a novel non-competitive NMDA antagonist, on motor and memory functions after closed head injury in the rat. *Brain Research*, 674(1), 55–62. [https://doi.org/10.1016/0006-8993\(94\)01433-I](https://doi.org/10.1016/0006-8993(94)01433-I)

Simon, G., Drori, J., & Cohen, M. (1967). Mechanism of conversion of aspartate into glutamate in cerebral-cortex slices. *Biochemical Journal*, 102(1), 153–162.

<https://doi.org/10.1042/bj1020153>

Singh, K., Trivedi, R., Haridas, S., Manda, K., & Khushu, S. (2016). Study of neurometabolic and behavioral alterations in rodent model of mild traumatic brain injury: A pilot study: Neurometabolic and behavioral alterations in rodent model of mTBI. *NMR in Biomedicine*, 29(12), 1748–1758. <https://doi.org/10.1002/nbm.3627>

Sperringer, Justin E., Adele Addington, and Susan M. Hutson. "Branched-chain amino acids and brain metabolism." *Neurochemical research* 42, no. 6 (2017): 1697-1709.

Stern, R. A., Riley, D. O., Daneshvar, D. H., Nowinski, C. J., Cantu, R. C., & McKee, A. C. (2011). Long-term Consequences of Repetitive Brain Trauma: Chronic Traumatic Encephalopathy. *PM&R*, 3(10 SUPPL. 2), S460–S467. <https://doi.org/10.1016/j.pmrj.2011.08.008>

Szczygielski, J., Hubertus, V., Kruchten, E., Müller, A., Albrecht, L. F., Schwerdtfeger, K., & Oertel, J. (2023). Prolonged course of brain edema and neurological recovery in a translational model of decompressive craniectomy after closed head injury in mice. *Frontiers in Neurology*, 14, 1308683. <https://doi.org/10.3389/fneur.2023.1308683>

Szczygielski, J., Mautes, A. E., Müller, A., Sippl, C., Glameanu, C., Schwerdtfeger, K., Steudel, W.-I., & Oertel, J. (2016). Decompressive Craniectomy Increases Brain Lesion Volume and Exacerbates Functional Impairment in Closed Head Injury in Mice. *Journal of Neurotrauma*, 33(1), 122–131. <https://doi.org/10.1089/neu.2014.3835>

Tepper, J. M., & Koós, T. (2016). GABAergic Interneurons of the Striatum. In *Handbook of Behavioral Neuroscience* (Vol. 24, pp. 157–178). Elsevier. <https://doi.org/10.1016/B978-0-12-802206-1.00008-8>

Tiwari, V., Ambadipudi, S., & Patel, A. B. (2013). Glutamatergic and GABAergic TCA Cycle and Neurotransmitter Cycling Fluxes in Different Regions of Mouse Brain. *Journal of Cerebral Blood Flow & Metabolism*, 33(10), 1523–1531. <https://doi.org/10.1038/jcbfm.2013.114>

Ulus, I. H., Wurtman, R. J., Mauron, C., & Blusztajn, J. K. (1989). Choline increases acetylcholine release and protects against the stimulation-induced decrease in phosphatide levels within membranes of rat corpus striatum. *Brain Research*, 484(1–2), 217–227.
[https://doi.org/10.1016/0006-8993\(89\)90364-8](https://doi.org/10.1016/0006-8993(89)90364-8)

Vagozzi, R., Tavazzi, B., Signoretti, S., Amorini, A. M., Belli, A., Cimatti, M., Delfini, R., Di Pietro, V., Finocchiaro, A., & Lazzarino, G. (2007). TEMPORAL WINDOW OF METABOLIC BRAIN VULNERABILITY TO CONCUSSIONS: MITOCHONDRIAL-RELATED IMPAIRMENT—PART I. *Neurosurgery*, 61(2), 379–389.
<https://doi.org/10.1227/01.NEU.0000280002.41696.D8>

Viant, M. R., Lyeth, B. G., Miller, M. G., & Berman, R. F. (2005). An NMR metabolomic investigation of early metabolic disturbances following traumatic brain injury in a mammalian model. *NMR in Biomedicine*, 18(8), 507–516. <https://doi.org/10.1002/nbm.980>

Wang, L.-N., Gu, X.-R., Si, N., Wang, H.-J., Zhou, Y.-Y., Bian, B.-L., & Zhao, H.-Y. (2022). [Biomarkers related to cognitive dysfunction in APP/PS1 mice based on non-targeted metabonomics and intervention mechanism of Huanglian Jiedu Decoction]. *Zhongguo Zhong Yao Za Zhi = Zhongguo Zhongyao Zazhi = China Journal of Chinese Materia Medica*, 47(22), 6117–6126. <https://doi.org/10.19540/j.cnki.cjcm.20220705.401>

Wang, X., Gao, X., Michalski, S., Zhao, S., & Chen, J. (2016). Traumatic Brain Injury Severity Affects Neurogenesis in Adult Mouse Hippocampus. *Journal of Neurotrauma*, 33(8), 721–733. <https://doi.org/10.1089/neu.2015.4097>

Xu, X.-J., Yang, M.-S., Zhang, B., Niu, F., Dong, J.-Q., & Liu, B.-Y. (2021). Glucose metabolism: A link between traumatic brain injury and Alzheimer's disease. *Chinese Journal of Traumatology = Zhonghua Chuang Shang Za Zhi*, 24(1), 5–10. <https://doi.org/10.1016/j.cjtee.2020.10.001>

Yarnell, A. M., Barry, E. S., Mountney, A., Shear, D., Tortella, F., & Grunberg, N. E. (2016). The Revised Neurobehavioral Severity Scale (NSS-R) for Rodents. *Current Protocols in Neuroscience*, 75(1). <https://doi.org/10.1002/cpns.10>

Zhang, C., Chen, R.-X., Zhang, Y., Wang, J., Liu, F.-Y., Cai, J., Liao, F.-F., Xu, F.-Q., Yi, M., & Wan, Y. (2017). Reduced GABAergic transmission in the ventrobasal thalamus contributes to thermal hyperalgesia in chronic inflammatory pain. *Scientific Reports*, 7(1), 41439. <https://doi.org/10.1038/srep41439>

Zheng, F., Zhou, Y., Feng, D., Li, P., Tang, T., Luo, J., & Wang, Y. (2020). Metabolomics analysis of the hippocampus in a rat model of traumatic brain injury during the acute phase. *Brain and Behavior*, 10(2). <https://doi.org/10.1002/brb3.1520>

Zheng, F., Zhou, Y.-T., Li, P.-F., Hu, E., Li, T., Tang, T., Luo, J.-K., Zhang, W., Ding, C.-S., & Wang, Y. (2020). Metabolomics Analysis of Hippocampus and Cortex in a Rat Model of Traumatic Brain Injury in the Subacute Phase. *Frontiers in Neuroscience*, 14, 876. <https://doi.org/10.3389/fnins.2020.00876>

Global, regional, and national burden of traumatic brain injury and spinal cord injury, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016

James, Spencer L et al. The Lancet Neurology, Volume 18, Issue 1, 56 – 87 Mitochondria Are Excitable Organelles Capable of Generating and Conveying Electrical and Calcium Signals Ichas, François et al. Cell, Volume 89, Issue 7, 1145 – 1153 Aspects of energy-linked calcium accumulation by rat heart mitochondria. Jacobus, WE et al. Journal of Biological Chemistry, Volume 250, Issue 19, 7863 – 7870 Development of the blood-brain barrier Risau, Werner et al. Trends in Neurosciences, Volume 13, Issue 5, 174 – 178.

Eliasson, Camilla et al Journal of Biological Chemistry, Volume 274, Issue 34, 23996 - 24006)

Chen, Xiao-Han, Robert Siman, Akira Iwata, David F. Meaney, John Q. Trojanowski, and Douglas H. Smith. "Long-term accumulation of amyloid- β , β -secretase, presenilin-1, and caspase-3 in damaged axons following brain trauma." The American journal of pathology 165, no. 2 (2004): 357-371.

Washington, Patricia M., Nicholas Morffy, Maia Parsadanian, David N. Zapple, and Mark P. Burns. "Experimental traumatic brain injury induces rapid aggregation and oligomerization of amyloid-beta in an Alzheimer's disease mouse model." Journal of neurotrauma 31, no. 1 (2014): 125-134.

Xu, Su, Jiachen Zhuo, Jennifer Racz, Da Shi, Steven Roys, Gary Fiskum, and Rao Gullapalli. "Early microstructural and metabolic changes following controlled cortical impact injury in rat: a magnetic resonance imaging and spectroscopy study." Journal of neurotrauma 28, no. 10 (2011): 2091-2102.

Pandya, J.D., Leung, L.Y., Yang, X., Flerlage, W.J., Gilsdorf, J.S., Deng-Bryant, Y., & Shear, D.A. (2019). Comprehensive Profile of Acute Mitochondrial Dysfunction in a Preclinical Model of Severe Penetrating TBI. *Frontiers in Neurology*, 10.

Viant, M., Lyeth, B., Miller, M., & Berman, R. (2005). An NMR metabolomic investigation of early metabolic disturbances following traumatic brain injury in a mammalian model. *NMR in Biomedicine*, 18. <https://doi.org/10.1002/NBM.980>.

Jiang XB, Ohno K, Qian L, Tominaga B, Kuroiwa T, Nariai T, Hirakawa K. Changes in local cerebral blood flow, glucose utilization, and mitochondrial function following traumatic brain injury in rats. *Neurol Med Chir (Tokyo)*. 2000 Jan;40(1):16-28; discussion 28-9. doi: 10.2176/nmc.40.16. PMID: 10721252.

Petroff, Ognen AC. "Metabolic biopsy of the brain." In Molecular neurology, pp. 77-100. Academic Press, 2007.

Sowers JL, Sowers ML, Shavkunov AS, Hawkins BE, Wu P, DeWitt DS, Prough DS, Zhang K. Traumatic brain injury induces region-specific glutamate metabolism changes as measured by multiple mass spectrometry methods. *iScience*. 2021 Sep 9;24(10):103108. doi: 10.1016/j.isci.2021.103108. PMID: 34622161; PMCID: PMC8479783.

P. Kontro, Interactions of Taurine and Dopamine in the Striatum, in: R.J. Huxtable, F. Franconi, A. Giotti (Eds.), *Biol. Taur.*, Springer US, Boston, MA, 1987: pp. 347–355. https://doi.org/10.1007/978-1-4899-0405-8_37.

F. Briand, E.A. Nelson, O. Maximo, W.P. Armstrong, N.V. Kraguljac, A.C. Lahti, Hippocampal glutamate and hippocampus subfield volumes in antipsychotic-naive first episode psychosis subjects and relationships to duration of untreated psychosis, *Transl. Psychiatry*. 10 (2020) 137. <https://doi.org/10.1038/s41398-020-0812-z>.

C.-S. Piao, A.L. Holloway, S. Hong-Routson, M.S. Wainwright, Depression following traumatic brain injury in mice is associated with down-regulation of hippocampal astrocyte glutamate transporters by thrombin, *J. Cereb. Blood Flow Metab. Off. J. Int. Soc. Cereb. Blood Flow Metab.* 39 (2019) 58–73. <https://doi.org/10.1177/0271678X17742792>.

J.R. Moffett, P. Arun, P.S. Ariyannur, A.M.A. Namboodiri, N-Acetylaspartate reductions in brain injury: impact on post-injury neuroenergetics, lipid synthesis, and protein acetylation, *Front. Neuroenergetics*. 5 (2013) 11. <https://doi.org/10.3389/fnene.2013.00011>.

M. Akram, Citric acid cycle and role of its intermediates in metabolism, *Cell Biochem. Biophys.* 68 (2014) 475–478. <https://doi.org/10.1007/s12013-013-9750-1>.

V. Di Pietro, A.M. Amorini, B. Tavazzi, R. Vagozzi, A. Logan, G. Lazzarino, S. Signoretti, G. Lazzarino, A. Belli, The Molecular Mechanisms Affecting N-Acetylaspartate Homeostasis Following Experimental Graded Traumatic Brain Injury, *Mol. Med.* 20 (2014) 147–157. <https://doi.org/10.2119/molmed.2013.00153>.

M. Kumari, P. Arora, P. Sharma, Y. Hasija, P. Rana, M.M. D'souza, N. Chandra, R. Trivedi, Acute metabolic alterations in the hippocampus are associated with decreased acetylation after blast induced TBI, *Metabolomics*. 19 (2023) 5. <https://doi.org/10.1007/s11306-022-01970-z>.

Littlewood-Evans A, Sarret S, Apfel V, Loesle P, Dawson J, Zhang J, Muller A, Tigani B, Kneuer R, Patel S, Valeaux S, Gommermann N, Rubic-Schneider T, Junt T, Carballido JM. GPR91 senses extracellular succinate released from inflammatory macrophages and exacerbates rheumatoid arthritis. *J Exp Med.* 2016;213:1655–1662.

Mills EL, Harmon C, Jedrychowski MP, Xiao H, Garrity R, Tran NV, Bradshaw GA, Fu A, Szpyt J, Reddy A, Prendeville H, Danial NN, Gygi SP, Lynch L, Chouchani ET. UCP1 governs liver extracellular succinate and inflammatory pathogenesis. *Nat Metab.* 2021;3:604–617.

Gibson GE, Xu H, Chen HL, Chen W, Denton TT, Zhang S. Alpha-ketoglutarate dehydrogenase complex-dependent succinylation of proteins in neurons and neuronal cell lines. *J Neurochem.* 2015;134:86–96.

Chen H, Xu H, Potash S, Starkov A, Belousov VV, Bilan DS, Denton TT, Gibson GE. Mild metabolic perturbations alter succinylation of mitochondrial proteins. *J Neurosci Res.* 2017;95:2244–2252.

Avram VF, Chamkha I, Åsander-Frostner E, Ehinger JK, Timar RZ, Hansson MJ, Muntean DM, Elmér E. Cell-permeable succinate rescues mitochondrial respiration in cellular models of statin toxicity. *Int J Mol Sci.* 2021;22:424.

Ehinger JK, Piel S, Ford R, Karlsson M, Sjövall F, Frostner E, Morota S, Taylor RW, Turnbull DM, Cornell C, Moss SJ, Metzsch C, Hansson MJ, Fliri H, Elmér E. Cell-permeable succinate prodrugs bypass mitochondrial complex I deficiency. *Nat Commun.* 2016;7:12317.

Ives SJ, Zaleski KS, Slocum C, Escudero D, Sheridan C, Legesse S, Vidal K, Lagalwar S, Reynolds TH. The effect of succinic acid on the metabolic profile in high-fat diet-induced obesity and insulin resistance. *Physiol Rep.* 2020;8:e14630.

Zhang Y, Zhang M, Zhu W, Yu J, Wang Q, Zhang J, Cui Y, Pan X, Gao X, Sun H. Succinate accumulation induces mitochondrial reactive oxygen species generation and promotes status epilepticus in the kainic acid rat model. *Redox Biol.* 2020;28:101365.

Huang LY, Ma JY, Song JX, Xu JJ, Hong R, Fan HD, Cai H, Wang W, Wang YL, Hu ZL, Shen JG, Qi SH. Ischemic accumulation of succinate induces Cdc42 succinylation and inhibits neural stem cell proliferation after cerebral ischemia/reperfusion. *Neural Regen Res.* 2023 May;18(5):1040-1045. doi: 10.4103/1673-5374.355821. PMID: 36254990; PMCID: PMC9827777.

Yudkoff, Marc. "Brain metabolism of branched-chain amino acids." *Glia* 21, no. 1 (1997): 92-98.

Hoffman, J. R., Zuckerman, A., Ram, O., Sadot, O., Stout, J. R., Ostfeld, I., & Cohen, H. (2017). Behavioral and inflammatory response in animals exposed to a low-pressure blast wave and supplemented with β -alanine. *Amino Acids*, 49(5), 871–886. <https://doi.org/10.1007/s00726-017-2383-8>

Hoge, C. W., McGurk, D., Thomas, J. L., Cox, A. L., Engel, C. C., & Castro, C. A. (2015). Mild Traumatic Brain Injury in U.S. Soldiers Returning from Iraq Charles. *New England Journal of Medicine*, 358(5), 453–463.

Kamnaksh, A., Ahmed, F., Kovesdi, E., Barry, E. S., Grunberg, N. E., Long, J. B., & Agoston, D. V. (2014). Molecular mechanisms of increased cerebral vulnerability after repeated mild blast-induced traumatic brain injury. *Translational Proteomics*, 3, 22–37. <https://doi.org/10.1016/j.trprot.2013.11.001>

Karagezyan, K.G., Ovsepyan, L.M. (1975). Role of ethanolamine, phosphoethanolamine, and phosphatidylethanolamine in oxidative phosphorylation of albino rat brain mitochondria. *Bull Exp Biol Med* 80, 1195–1196. <https://doi.org/10.1007/BF00833158>

Katada, S., Imhof, A., & Sassone-Corsi, P. (2012). Connecting Threads: Epigenetics and Metabolism. *Cell*, 148(1–2), 24–28. <https://doi.org/10.1016/j.cell.2012.01.001>

Lin, A. P., Shic, F., Enriquez, C., & Ross, B. D. (2003). Reduced glutamate neurotransmission in patients with Alzheimer's disease?an in vivo ^{13}C magnetic resonance spectroscopy study. *MAGMA Magnetic Resonance Materials in Physics, Biology and Medicine*, 16(1), 29–42. <https://doi.org/10.1007/s10334-003-0004-x>

Mishra, V., Skotak, M., Schuetz, H., Heller, A., Haorah, J., & Chandra, N. (2016). Primary blast causes mild, moderate, severe and lethal TBI with increasing blast overpressures: Experimental rat injury model. *Scientific Reports*, 6(1), 26992. <https://doi.org/10.1038/srep26992>

Patet, C., Suys, T., Bloch, J., Daniel, R. T., Pellerin, L., Magistretti, P. J., et al. (2015). Neuroenergetic response to prolonged cerebral glucose depletion after severe brain injury and the role of lactate. *Critical Care*, 19(1), 1–27.

Peskind, E. R., Petrie, E. C., Cross, D. J., Pagulayan, K., McCraw, K., Hoff, D., et al. (2011). Cerebrocerebellar hypometabolism associated with repetitive blast exposure mild traumatic brain injury in 12 Iraq war Veterans with persistent post-concussive symptoms. *NeuroImage*, 54(SUPPL. 1), S76–S82. <https://doi.org/10.1016/j.neuroimage.2010.04.008>

Prins, M., Alexander, D., Giza, C. C., & Hovda, D. A. (2013). Repeated Mild Traumatic Brain Injury: Mechanisms of Cerebral Vulnerability. *The Journal of neurotrauma*, 30(1), 30–38. <https://doi.org/10.1089/neu.2012.2399>

Rae, C., Hare, N., Bubb, W. A., McEwan, S. R., Bröer, A., McQuillan, J. A., et al. (2003). Inhibition of glutamine transport depletes glutamate and GABA neurotransmitter pools: Further evidence for metabolic compartmentation. *Journal of Neurochemistry*, 85(2), 503–514. <https://doi.org/10.1046/j.1471-4159.2003.01713.x>

Rana, P., Rama Rao, K. V., Ravula, A., Trivedi, R., D’Souza, M., Singh, A. K., et al. (2020). Oxidative stress contributes to cerebral metabolomic profile changes in animal model of blast-induced traumatic brain injury. *Metabolomics*, 16(3), 1–12. <https://doi.org/10.1007/s11306-020-1649-4>

Reichelt, K. L., & Kvamme, E. (1967). Acetylated and Peptide Bound Glutamate and Aspartate in Brain. *Journal of Neurochemistry*, 14(10), 987–996. <https://doi.org/10.1111/j.1471-4159.1967.tb09510.x>

Roberto, V., Signoretti, S., Tavazzi, B., Floris, R., Ludovici, A., Marziali, S., & Tarascio, G. (2008). Temporal window of metabolic brain vulnerability to concussion: A pilot ¹Hmagnetic resonance spectroscopic study in concussed athletes - Part III: Commentary. *Neurosurgery*, 62(6), 1286–1296. <https://doi.org/10.1227/01.NEU.0000316421.58568.AD>

Shimomura, Y., & Kitaura, Y. (2018). Physiological and pathological roles of branched-chain amino acids in the regulation of protein and energy metabolism and neurological functions. *Pharmacological Research*, 133(May), 215–217. <https://doi.org/10.1016/j.phrs.2018.05.014>

Stern, R. A., Riley, D. O., Daneshvar, D. H., Nowinski, C. J., Cantu, R. C., & McKee, A. C. (2011). Long-term Consequences of Repetitive Brain Trauma: Chronic Traumatic

Encephalopathy. PM&R, 3(10 SUPPL. 2), S460–S467.
<https://doi.org/10.1016/j.pmrj.2011.08.008>

Subramaniam, A., Agarwal, S., Kalmady, S., Shivakumar, V., Chhabra, H., Bose, A., et al. (2015). Effect of transcranial direct current stimulation on prefrontal inhibition in schizophrenia patients with persistent auditory hallucinations: A study on antisaccade task performance. Indian Journal of Psychological Medicine, 37(4), 419–422. <https://doi.org/10.4103/0253-7176.168584>

Takahashi, H., McCaffery, J. M., Irizarry, R. A., & Boeke, J. D. (2006). Nucleocytosolic Acetyl-Coenzyme A Synthetase Is Required for Histone Acetylation and Global Transcription. Molecular Cell, 23(2), 207–217.
<https://doi.org/10.1016/j.molcel.2006.05.040>

Truckenmiller, M. E., Namboodiri, M. A. A., Brownstein, M. J., & Neale, J. H. (1985). NAcetylation of L-Aspartate in the Nervous System: Differential Distribution of a Specific Enzyme. Journal of Neurochemistry, 45(5), 1658–1662. <https://doi.org/10.1111/j.1471-4159.1985.tb07240.x>

Tyagi, Ritu, Poonam Rana, Ahmad Raza Khan, Deepak Bhatnagar, M. Memita Devi, Shubhra Chaturvedi, Rajendra P. Tripathi, and Subash Khushu(2011). "Study of acute biochemical effects of thallium toxicity in mouse urine by NMR spectroscopy." Journal of applied toxicology 31, no. 7 663-670.

Van Hall, G., Strømstad, M., Rasmussen, P., Jans, Ø., Zaar, M., Gam, C., et al. (2009). Blood lactate is an important energy source for the human brain. Journal of Cerebral Blood Flow and Metabolism, 29(6), 1121–1129. <https://doi.org/10.1038/jcbfm.2009.35> Vazquez, A., Kamphorst, J. J., Markert, E. K., Schug, Z. T., Tardito, S., & Gottlieb, E. (2016). Cancer metabolism at a glance. Journal of Cell Science, 129(18), 3367–3373. <https://doi.org/10.1242/jcs.181016>

Weinhold, B. (2006). Epigenetic S. Environmental Health, 114(3), 160–167.

Wellen, K. E., Hatzivassiliou, G., Sachdeva, U. M., Bui, T. V., Cross, J. R., & Thompson, C. B. (2009). ATP-citrate lyase links cellular metabolism to histone acetylation. Science, 324(5930), 1076–1080. <https://doi.org/10.1126/science.1164097>

Xu, Su, Jiachen Zhuo, Jennifer Racz, Da Shi, Steven Roys, Gary Fiskum, and Rao Gullapalli. "Early microstructural and metabolic changes following controlled cortical impact injury in rat: a

magnetic resonance imaging and spectroscopy study." *Journal of neurotrauma* 28, no. 10 (2011): 2091-2102.

Xu, X., Cowan, M., Beraldo, F., Schranz, A., McCunn, P., Geremia, N., et al. (2021). Repetitive mild traumatic brain injury in mice triggers a slowly developing cascade of long-term and persistent behavioral deficits and pathological changes. *Acta Neuropathologica Communications*, 9(1), 1–27. <https://doi.org/10.1186/s40478-021-01161-2>

Zeisel, S. H. (1992). Choline: an important nutrient in brain development, liver function and carcinogenesis. *Journal of the American College of Nutrition*, 11(5), 473–481. <https://doi.org/10.1080/07315724.1992.10718251>

Zeisel, S. H., Mar, M.-H., Howe, J. C., & Holden, J. M. (2003). Concentrations of Choline-Containing Compounds and Betaine in Common Foods. *The Journal of Nutrition*, 133(5), 1302–1307. <https://doi.org/10.1093/jn/133.5.1302>

Zhou, Y., & Danbolt, N. C. (2014). Glutamate as a neurotransmitter in the healthy brain. *Journal of Neural Transmission*, 121(8), 799–817. <https://doi.org/10.1007/s00702-0141180-8>

Publications from thesis

- **Kumari, Megha**, Palkin Arora, Priyanka Sharma, Yasha Hasija, Poonam Rana, Maria M. D'souza, Namas Chandra, and Richa Trivedi. "Acute metabolic alterations in the hippocampus are associated with decreased acetylation after blast induced TBI." *Metabolomics* 19, no. 1 (2023): 5.
- **Kumari, Megha**, Yasha Hasija, and Richa Trivedi. "Acute and sub-acute metabolic change in different brain regions induced by moderate blunt traumatic brain injury." *NeuroReport* 35, no. 2 (2024): 75-80.

Other publications

- Arora P, Singh K, **Kumari M**, Trivedi R. Temporal profile of serum metabolites and inflammation following closed head injury in rats is associated with HPA axis hyperactivity. *Metabolomics*. 2022;18(5):28. Published 2022 Apr 29. doi:10.1007/s11306-022-01886-8.
- Palkin Arora, Richa Trivedi, **Megha Kumari**, Kavita Singh, Rajat Sandhir, Maria M. D'Souza, and Poonam Rana. "Altered DTI scalars in the hippocampus are associated with morphological and structural changes after traumatic brain injury." *Brain Structure and Function* (2024): 1-11.
- Palkin Arora, **Megha Kumari**, Kavita Singh, Poonam Rana, Rajat Sandhir, Richa Trivedi Hyperacute metabolic alterations in TBI induced rats are associated with hypothalamic structural impairment and gut microbiome changes, *Journal of Neuroscience Research*, Submitted

Book chapter

- **Kumari, Megha**, Palkin Arora, and Richa Trivedi. "Epigenetic and metabolic changes in traumatic brain injury." In *Epigenetics and Metabolomics*, pp. 97-106. Academic Press, 2021.